Peoples ’ Democratic Republic of Algeria

Ministry of Higher Education and Scientific Research

University of Mentouri- Constantine 1

Faculty of Exact Sciences

Department of Mathematics



Acknowledgments

In the name of ALLAH, the Most Gracious and the Most Merciful

Alhamdulillah, all praises to ALLAH for the Strengths and His Blessing in completing this document.

I would like to thank all the mathematics module managers that I worked with them BOUCHEKOUF Messaouda, REKKAB Soraya, ARAR Nouria and SMAKDJI Mohamed EL-Hadi. They have allowed me to benefit from their experiences and for all the information, advice and also the confidence they gave me.

My sincere thanks also go to Prof. DALAH Mohamed and Prof. KHELLAF Hassane. Without their precious expertise and advises it would not be possible to publish this document.


To understand the Universe, you must understand the language in which it's written,
the language of Mathematics.
Galileo Galilei
Do not worry about your difficulties in Mathematics. I can assure you mine are still greater.
Albert Einstein

Introduction

"To understand the Universe, you must understand the language in which it's written, the language of Mathematics" Galileo Galilei

This is a Galileo's quotation which has been proved over time and with the development of the mathematical theories. Thus we can describe more and more phenomena of the universe whether they are physical, biological, ecological, economic ... through different laws and mathematical entities.

Dear student, if you want to become an engineer or a specialist in any of the university scientific fields, you must have a good achievement in mathematics in order to pass to the other sciences and to achieve higher academic degrees.

This handout in General Algebra, Analysis and Linear Algebra includes chapters that cover the vocabulary of the Mathematics 1 curriculum which a first year student in Science of Technology (ST) specialty in the first semester. This scientific production can be used as a reference for undergraduates in the first year Mathematics and other Experimental Sciences majors as well.

In fact, we have noticed the absence of references on the ministerial curriculum in the mathematics module in English, for this reason we provide this work for Algerian and foreign students as an attempt to overcome the difficulties facing them in English, the contemporary language of science, and provide them with the common terms to stimulate their educational activity and scientific research in this language, as it is the language of the most prominent scientific web sites

and the language of business and international conferences.

The topics were formulated appropriately in a simple linguistic style, taking into account the necessary accuracy and easiness. All the concepts were followed by examples and solved exercises.

1. Logic and mathematical reasoning

The first year of university is an essential stage on which the success or the failure of the coming years depends, as it adopts the integrity of thinking of the student with mathematical logic which is the basis of mathematics and is the subject of ***the first chapter*** *of the program. Through it, they learn the logical steps of solving a problem with the help of patterns of proof. Logic is used also in the field of working with devices such as computers and electrical circuits.*

2. Sets, Relations and Applications

The ***second chapter*** *is devoted to the set theory, starting from the sets concepts and belonging in the sets, operations and the relations that gave rise to the concept of applications and functions. Thus, this fundamental theory is particularly appreciated in physics, for example in the study of electrical circuits and in chemistry, in the study of atomic orbital's, and it also plays a major role in the design and construction of electronic computers, as it is largely involved in information organization, management techniques and market studies.*

3. *Real-valued functions of a real variable*

*The **third chapter** presents the real function. Which is an ancient mathematical concept that appeared with the first human civilizations, The Babylonians who have the credit for its appearance, used numerical tables to highlight the relationship between two quantities, for example to make calculations (tables of squares, square roots, cubes and cubic roots) or to compile the astronomical calendar for the sun, moon or other planets.*

The concept of function is a universal concept that is largely founded in all practical scientific disciplines: mathematics, physics, biology, technology and even in the human sciences...

In this chapter, we will study the limits, continuity, the Intermediate Value Theorem, differentiability of real functions, the Mean Value Theorem and L'Hospital's Rule which is used to get us out of sticky situations with indeterminate limit forms.

4. *Finite Expansions*

Taylor's series expansion formula is a new concept for first-year university students. It is used to approximate a differentiable function locally (that is, near a specific point) with polynomial which can facilitate many of the calculations performed by mathematicians.

*The idea of approximation was generalized even for the function which is not locally defined or not differentiable to a polynomial. Finite expansions are the subject of **the fourth chapter**.*

They are useful tools for mathematic analysis and physics. In particular, they provide a good way to solve some indeterminate forms of limits.

5. *Vector Space and Linear Maps*

Dear student, you have studied how to solve linear equations system with 2 unknowns and the question now is, what must you do if the number of the unknowns is more than 2?

This problem was discussed by René Descartes, who has proposed a new algebraic structure; the vector spaces which is the content of the ***last chapter****.*

The Mathematical Modelling of Natural Phenomena is done with vector spaces and these models can be used to make effective calculations.

As for nonlinear systems that cannot be modelled by linear algebra and vector spaces, they are usually used to deal with first-order approximations.

We focus our research on those elementary concepts, which appeared to be essential: linear combination, independence-dependence, generating sets, bases, and rank-dimension, linear map.

Appendix

At the end, we presented a ***summary of the courses translated into Arabic in the appendices A,B,C,D,E*** *to facilitate the understanding of the English mathematical terms.*

In conclusion, we hope that this publication, in its content, organization and methodology, will be an effective support for our students, a flexible support for professors, and a contribution to enriching the Algerian library.

I found a great pleasure in writing in English. I hope that this fun will accompany students who focus their interests on seeking knowledge and ascending to higher levels.

Doctor KESMIA Mounira

The future is for those who believe in the beauty of their dreams .

Success comes from GOD

مقدمة

" لفهم الكون ، يجب أن تفهم اللغة التي كُتب بها ، لغة الرياضيات" غاليلي غاليليو

هذه المقولة لغاليلي تؤول إلى التحقق بمرور الوقت ومع تطور النظريات الرياضياتية. وبالتالي نتمكن من وصف المزيد والمزيد من ظواهر الكون سواء كانت فيزيائية أو بيولوجية أو بيئية أو اقتصادية ... من خلال قوانين وكيانات رياضياتية مختلفة.

فإذا أردت عزيزي الطالب أن تصبح مهندسا أو متخصصا في أي من المواد العلمية الجامعية لابد من تحصيل رياضياتي جيد للعبور إلى العلوم التجريبية وإلى تحقيق مراتب علمية عالية .

إن هذه المطبوعة في الجبر العام والتحليل والجبر الخطي تتضمن فصول تغطي مفردات منهاج رياضيات 1 وفقا للمنهاج الوزاري المقرر الذي يدرسه طلاب السنة الأولى علوم التكنولوجيا (ST) في السداسي الأول وكذلك طلاب شعبة الرياضيات وباقي العلوم التجريبية.

ونظرا لملاحظتنا غياب مراجع باللغة الإنجليزية للمنهاج الوزاري في مقياس الرياضيات, ارتأينا وضع هذا العمل المتواضع بين أيدي طلبتنا الأعزاء من أبناء الوطن و الأجانب, محاولة منا لتذليل عقبة لغة العلم المعاصرة وتزويد طلابنا بالمصطلحات المتداولة لتحفيز النشاط التعليمي والبحثي لهم بهذه اللغة, فهي لغة أبرز المواقع العلمية الموجهة عبر الإنترنت, و لغة الأعمال والملتقيات الدولية.

هذه المطبوعة مهيكلة في فصول بصورة مناسبة ومصاغة بأسلوب لغوي بسيط مع مراعاة الدقة اللازمة وسهلة الاستعمال بحيث اتبعت الفقرات بأمثلة وتمارين محلولة.

1. المنطق الرياضي وأنماط البرهان

تعتبر السنة أولى جامعي مرحلة أساسية يتوقف عليها نجاح أو فشل السنوات المقبلة, إذ أنه خلالها تبنى سلامة التفكير عند الطالب بالمنطق الرياضي الذي يعتبر أساس الرياضيات وهو موضوع الفصل الأول من المنهاج. من خلاله يتعلم الخطوات المنطقية لحل مسألة بالاستعانة بأنماط البرهان ويساعد على اكتساب مهارات التفكير والتحليل المنطقي والاستنتاج العلمي. كما يستخدم المنطق في مجال العمل مع الأجهزة مثل أجهزة الكمبيوتر والدوائر الكهربائية.

2. المجموعات, العلاقات والتطبيقات

أما الفصل الثاني فهو مكرس لنظرية المجموعات , بدءًا من مفاهيم المجموعة والانتماء, العمليات في المجموعات والعلاقات بين عناصر نفس المجموعة والعلاقة من مجموعة نحو أخرى التي نشأ عنها مفهوم التطبيقات والدوال التي وجدت تطبيقات عديدة مثلا في الفيزياء دراسة الدارات الكهربائية و في الكيمياء دراسة المدارات الذرية, كما تلعب دورًا كبيرًا في تصميم وبناء أجهزة الكمبيوتر الإلكترونية, و تشارك كذلك إلى حد كبير في تنظيم المعلومات وتقنيات الإدارة ودراسات السوق.

3. الدوال الحقيقية بمتغير حقيقي واحد

يعرض الفصل الثالث الدالة الحقيقية, هذا المفهوم القديم قدم الرياضيات الذي ظهر مع الحضارات الإنسانية الأولى, فالبابليون لهم الفضل في ظهوره, الذين استخدموا الجداول العددية لإبراز العلاقة بين مقدارين, مثلا لإجراء الحسابات (جداول المربعات والجذور التربيعية والمكعبات والجذور التكعيبية) أو لتجميع التقويم الفلكي للشمس والقمر أو الكواكب الأخرى . يعد مفهوم الدالة مفهومًا عالميًا إلى حد كبير موجود في جميع التخصصات العلمية عمليًا: الرياضيات والفيزياء و البيولوجيا والتكنولوجيا ولكن أيضًا في العلوم الإنسانية ...

في هذا الفصل نتطرق إلى النهايات , الاستمرار , نظرية القيم المتوسطة, الاشتقاق, نظرية التزايدات المنتهية و قاعدة لوبيتال التي تمكننا من إزالة بعض حالات عدم التعيين وحساب النهايات للدوال الحقيقية.

4. النشر المنتهي أو النشر المحدود

نشر تايلور يعد مفهوما جديدا نقدمه لطلاب السنة أولى جامعي, والذي ماهو إلا تقريبا لدالة قابلة الاشتقاق من رتب عليا في جوار نقطة معينة, بكثير حدود من شأنه أن يسهل الكثير من الحسابات التي يقوم بها الرياضيون.

تم تعميم فكرة النشر إلى تقريب دالة في الحالة العامة حتى وإن كانت هذه الدالة غير معرفة أو غير قابلة الاشتقاق في جوار نقطة معينة, إلى كثير حدود يسمى النشر المنتهي أو النشر المحدود وهو موضوع الفصل الرابع, بحيث يعتبر أداة فعالة في التحليل الرياضي والفيزياء. كما يسمح النشر المحدود برفع حالات عدم التعيين في العبارات التي نجدها مثلا في دراسة الدوال أو البحث عن النهايات.

5. الفضاء الشعاعي و التطبيقات الخطية

لقد درست عزيزي الطالب كيفية حل جمل معادلات خطية ذات مجهولين, والسؤال المطروح الآن, فما هو الحل في حالة ما إذا كان عدد المجاهيل أكبر من ذلك؟

هذه الإشكالية جعلت رينيه ديكارت يفكر في بنية جبرية جديدة فكانت الفضاءات الشعاعية محتوى الفصل الأخير التي وجدت طريقا إلى باقي العلوم فهي وسيلة ممتازة لنمذجة الظواهر الطبيعية ويمكن استخدام هذه النماذج لإجراء حسابات فعالة, وفيما يتعلق بالأنظمة غير الخطية التي لا يمكن نمذجتها عن طريق الجبر الخطي و الفضاءات الشعاعية, فإنه يتم استخدامها عادةً للتعامل مع التقديرات التقريبية من

الدرجة الأولى. ونتناول في هذا الفصل التعريف بهذه البنية, حيث نقدم المفاهيم الأولية كالمزج الخطي, الجملة المولدة, الاستقلال والارتباط الخطي, الأسس, الأبعاد, التطبيقات الخطية و الرتبة.

الملحقات

في النهاية، قدمنا ملخصًا للفصول مترجمة إلى العربية في الملحقات لتسهيل فهم المصطلحات الرياضياتية بالانجليزية.

وفي الختام نرجو أن تكون مطبوعتنا هذه في محتواها وتنظيمها ومنهجيتها سندا فعالا لطلبتنا الأعزاء وسندا مرنا للأساتذة وإسهاما في إثراء المكتبة الجزائرية.

لقد وجدنا المتعة المتزايدة عند تأليفها بالانجليزية . ونتمنى أن ترافق هذه المتعة, الطلبة الذين يركزون اهتماماتهم على طلب العلم والارتقاء إلى الدرجات العلى.

الدكتورة قسمية منيرة

المستقبل ملك لأولئك الذين

يؤمنون بجمال أحلامهم

بتوفيق من الله.

Contents

1 Logic and mathematical reasoning **18**

1.1 Propositional Logic 18

1.2 Truth table 19

1.3 Logical Connectors 20

1.3.1 Conjunction 20

1.3.2 Disjunction 20

1.3.3 Implication 20

1.3.4 Equivalence 23

1.4 Predicates and Quantifiers 30

1.4.1 Nested Quantifiers 32

1.5 Methods of Proof 34

1.5.1 Direct Method 35

1.5.2 Proof by Contrapositive 35

1.5.3 Proof by Contradiction (Absurd) 36

1.5.4 Proof by Counter-Example 37

1.5.5 Proof by Cases Disjunction 38

1.5.6 Proof by Mathematical Induction 39

2 Sets, Relations and Applications **41**

2.1 Set Theory 41

2.1.1 Relationships between elements and parts of a set 41

2.1.2 Set Operations 46

2.2 Relations 57

2.2.1 Binary Relations 58

2.2.2 Equivalence Relation 59

2.2.3 Equivalences and Partitions 61

2.2.4 Order Relation 62

2.3 Applications 64

2.3.1 Functional Relation 64

2.3.2 Equality - Extension - Restriction 66

2.3.3 Image and Inverse Image of a Subsets 67

2.3.4 Injective, Surjective and Bijective Applications 69

2.3.5 Examples of Applications 73

2.3.6 Operations on Applications 74

2.4 The Inverse Trigonometric Application 78

2.4.1 Arccosine Application 78

2.4.2 Arcsine Application 79

2.4.3 Arctangent Application 82

2.4.4 Arccotangent Application 83

3 Real functions of one real variable **86**

3.1 Elementary Functions 86

3.2 Limit of a Function 87

3.2.1 Limit of a function at a point 87

3.2.2 Limit of a function at infinity 90

3.2.3 Properties of the limit 91

3.2.4 Indeterminate Forms 92

3.3 Continuous Functions 94

3.3.1 Continuity of a function at a point 94

3.3.2 Continuity of a function in an interval 95

3.3.3 Continuous Extension at a point 96

3.3.4 The Intermediate Value Theorem 97

3.3.5 Continuity of composite functions 98

3.3.6 Continuity of the algebraic combinations of functions 98

3.4 Differentiability of Functions 99

3.4.1 Differentiability of a function at a point 99

3.4.2 Differentiability of a function in an interval 102

3.4.3 Derivative Table of Elementary Functions 102

3.4.4 Derivative of inverse function 103

3.4.5 Algebra of derivatives 103

3.4.6 Derivatives of composite functions 103

3.4.7 Derivative of some composite Functions 104

3.4.8 Lagrange's Mean Value Theorem 104

3.4.9 L'Hospital's Rule 104

3.4.10 Higher derivatives 107

4 Finite Expansions **109**

4.1 Approximating functions by Taylor polynomials 109

4.1.1 First degree Taylor polynomials 109

4.1.2 Mac-Laurin Polynomials 112

4.2 Finite expansions at zero 113

4.2.1 Basic finite expansions of elementary functions 114

4.3 Algebraic combinations of finite expansions 115

4.4 Composite of finite expansions 117

4.5 Finite expansions at a point 118

4.6 Finite expansions at Infinity 119

4.7 Using finite expansions to evaluate limits 120

5 Vector Space and Linear Maps 122

5.1 Vector Space 122

5.1.1 Subspaces of a vector space 124

5.1.2 Linear combinations 125

5.1.3 Linear independence and linear dependence 126

5.1.4 Generating sets 127

5.1.5 Bases of a vector space 128

5.1.6 Dimension of a vector space 129

5.2 Linear Maps 130

5.2.1 Linear maps and dimension 132

A Logic and mathematical reasoning translated 3

B Sets, Relations and Applications translated 9

C Real functions of one real variable translated 22

D Finite Expansions translated 29

E Vector Space and Linear Maps translated 35

FAMOUS ISLAMIC MATHEMATICIANS

Muhammad Ibn Musa Al-Khwarizmi

The basis of **Algebra** was found by Al Khwarizmi, He is also known as the person who introduced the Western world to the decimal numerical system, which is the most widely used number

Let's REMEMBER

Number Sets

In mathematics very often we study sets whose elements are the real numbers. Some special number sets which are frequently encountered are defined as follow.

- ♦ $\mathbb{N}$ is the set of **Natural numbers**: $\mathbb{N} = \{0, 1, 2, 3, ...\}$
- ♦ $\mathbb{Z}$ is the set of **Integers:** $\mathbb{Z} = \{....., -3, -2, -1, 0, 1, 2, 3, ...\}$
- ♦ D is the set of **Decimal numbers:** $D = \left\{\frac{p}{10^n}, p \in \mathbb{Z}, n \in \mathbb{N}\right\}$

Example: $1.234 = \frac{1234}{10^4}$ is a decimal number.

- ♦ $\mathbb{Q}$ is the set of **Rational numbers:** $\mathbb{Q} = \left\{\frac{p}{q}, p \in \mathbb{Z}, q \in \mathbb{Z}^*\right\}$.

Rational numbers are numbers that can be expressed as the quotient of two integers (ie a fraction) with a denominator that is not zero. Note that all terminating decimals or repeating decimals (or periodic decimal expansion) are rational numbers.

Examples:

1. $\frac{1}{2} = 0.5$ (terminating decimals).
2. $\frac{9}{7} = 1,285714\mathbf{285714}28571428... = 1,\overline{285714}$ (repeating decimals)

- ♦ $\mathbb{R}$ is the set of **Real numbers**, numbers that can be represented by any decimal expansion, limited or not.

Example: 123.101001000100001....,etc.

♦ $\mathbb{C}$ is the set of **Complex numbers** $\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}\}$

Recall that a complex number is formed by adding a real number to a real multiple of i, where $i = \sqrt{-1}$.

We have $\mathbb{N} \subset \mathbb{Z} \subset D \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$

♦ The set of **Even numbers** contains the elements $0, \pm 2, \pm 4, \pm 6, ...$ which are those of the form $2n$ for some integer n.

♦ The set of **Odd numbers** is the set of integers which are not even. Hence odd numbers are $\pm 1, \pm 3, \pm 5, ...$ which can be written as $2n + 1$ for some integer n.

Absolute value

For real numbers x we define the absolute value of x to be

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

Examples: $|-2| = 2$, $|\sqrt{2}| = \sqrt{2}$, and $|0| = 0$.

Properties:

1. $|x| \geq 0$
2. $\sqrt{x^2} = |x|$
3. $|x|^2 = x^2$
4. $|x \cdot y| = |x| \cdot |y|$
5. $|x + y| \leq |x| + |y|$

The greatest integer function

For real numbers x, the *greatest integer function* $[x]$ gives the greatest integer not greater than x.

Examples: $[3.14] = 3$, $[-3.14] = -4$, $[0.7] = 0$.

The general solution to the quadratic equations and inequalities

We only consider here the case $a > 0$ to which we can always reduce

$\Delta = b^2 - 4ac$ $(a > 0)$	Solution of the equation $ax^2 + bx + c = 0$	Solution of the inequality $ax^2 + bx + c > 0$	Solution of the inequality $ax^2 + bx + c < 0$
If $\Delta > 0$	$x_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $(x_1 < x_2)$	$(x < x_1) \vee (x > x_2)$	$(x_1 < x < x_2)$
If $\Delta = 0$	$x_1 = x_2 = \frac{-b}{2a}$	$\forall x \neq \frac{-b}{2a}$	no solution
If $\Delta < 0$	no solution	$\forall x \in \mathbb{R}$	no solution

Let's remember that:

1. $ax^2 + bx + c = a(x - x_1)(x - x_2)$.
2. $\begin{cases} x_1 + x_2 = \frac{-b}{a} \\ x_1 \cdot x_2 = \frac{c}{a} \end{cases}$

Identities and Trigonometric Formulas

1. $\cos x = \frac{adjacent}{hypotense}$, $\sin x = \frac{opposite}{hypotense}$
2. $\tan x = \frac{\sin x}{\cos x}$, $\cot x = \frac{1}{\tan x}$
3. $\cos x$ and $\sin x$ defined for all real numbers $x \in \mathbb{R}$
4. $\tan x$ defined for all $x \neq (2n+1)\frac{\pi}{2}$, where $n \in \mathbb{Z}$
5. $\cot x$ defined for all $x \neq n\pi$, where $n \in \mathbb{Z}$
6. $-1 \leq \cos x \leq 1$, $-1 \leq \sin x \leq 1$, $-\infty \leq \tan x \leq +\infty$, $-\infty \leq \cot x \leq +\infty$
7. $\sin x^2 + \cos x^2 = 1$

FAMOUS ISLAMIC MATHEMATICIANS



Abu Al-Hasan

Abu Al-Hasan was the first person to efficiently add up the first hundred numbers. He did a lot of work on the **number theory**, and came across theories based upon perfect numbers. He created theories linking **algebra** to **geometry**.

Chapter 1

Logic and mathematical reasoning

Logic is the hygiene the mathematician practices to keep his ideas healthy and strong

"Hermann Weyl"

Mathematical logic is a subfield of mathematics exploring the study of reasoning. It analyzes the structure of the methods and validity of mathematical deduction and proof.

The principles of logic can be attributed to ARISTOTLE (384–322 B.C.E.), who proposed as formal rules for correct reasoning. He searched to identify modes of reasoning that are valid by virtue of their structure, not their content. For example, "1 and 2 are numbers; therefore 2 is a number" and "$\sqrt{5}$ and π are natural numbers; therefore π is natural number" have the same structure ("A and B, therefore A"), and any reasoning made via this structure is logically valid. (In particular, the second example is logically sound.)

1.1 Propositional Logic

Definition (Proposition)

A proposition is a statement which has a truth value either true or false.

Notation: Variables are used to represent propositions. The most common variables used are p, q, and r.

Examples:

1. p :"2 is even", q :"2 + 2 = 4", r :"2 + 2 = 5" are popositions.

2. "$x + 2 = 2x$" is not a proposition.

Definition (Negation)

The negation of a proposition p is also called **not** p, and is denoted by $\bar{p}$.

Examples: Give the negation of the following statements.

1. If p: "2 is even" then $\bar{p}$: "2 is not even".

2. If p: "2 + 2 = 5" then $\bar{p}$: "2 + 2$\neq$ 5".

1.2 Truth table

Definition (Truth-value)

The **truth-value** is one of the two values, "true" (T) or "false" (F), that can be taken by a given logical formula in an interpretation (model) considered. Sometimes the truth value T is denoted in the literature by 1, and F by 0.

Definition (Truth table)

A **truth table** is a table showing the truth-value of a statement (typically a compound one) given the possible truth-values of the simple statements of which it is composed.

The truth values of a proposition, p, can be displayed in tabular form as follows:

p
1
0

1.3 Logical Connectors

1.3.1 Conjunction

Definition (Conjunction)

If p and q are two propositions then their **conjunction** is the proposition whose value is true only when both are true. A conjunction can also be written $p \wedge q$ which is read p **and** q.

Examples:

1. "A triangle has three sides and a square has four sides" is a conjunction

2. Let p : "$2 \leq 3$" and q : "$2^2 \leq 3^2$" , the proposition $p \wedge q$ is true.

1.3.2 Disjunction

Definition (Disjunction)

A compound statement of the form "p **or** q" is known as a disjunction and it is denoted by $p \vee q$. The **disjunction** of p and q has value false only when both are false.

Examples:

1. "An integer is a number which presents itself as a natural integer to which a positive **or** negative sign has been added indicating its position relative to 0 on an oriented axis ".

2. $\bar{p} \vee \bar{q}$: "$2 > 3$" $\vee$ "$2^2 > 3^2$" is a false proposition.

1.3.3 Implication

Definition (Implication)

A **conditional** statement of the form "**If** ... **then**..." is known as a **conditional** or an **implication**.

A conditional statement has two components: If p, then q. Statement p is called the antecedent (hypothesis, or premise) and statement q the consequent (or conclusion).

Alternative Phrasings of Conditionals

A conditional statement can be written a number of different, but equivalent, ways:

If p, then q.

p implies q.

q if p.

p only if q.

p is sufficient for q.

q is necessary for p.

It is denoted in symbols by: $p \Rightarrow q$.

The implication of p and q has value false only when p is true and q is false.

Examples:

1. "If a polygon has three sides, then it is a triangle" is a conditional statement.

2. "If $1 \leq 3$ then $1 + 1 \leq 3 + 1$" is a true implication.

3. "If π and $2 + 3i$ are real numbers then $2 + 3i$ is real number" is a true implication.

4. "If $2 + 3 = 5$ then $3 \times 2 + 3 \times 3 = 20$" is a false implication because when $x = 5$, $3x = 15$ and $15 \neq 20$.

5. "If $(-2)^2 = 4$ then $-2 = \sqrt{4}$" is a false implication because $\sqrt{(-2)^2} \neq -2$.

Definition (Converse of implication)

The converse of $p \Rightarrow q$ is the proposition $q \Rightarrow p$.

Example:

Let p : "x is a prime number different from 2" and q : "x is odd". One has $p \Rightarrow q$ but we do not have $q \Rightarrow p$.

Theorem 1. For all propositions p and q, the following statements are true.

1. $p \Rightarrow p \vee q$ and $q \Rightarrow p \vee q$

2. $p \wedge q \Rightarrow p$ and $p \wedge q \Rightarrow q$

Proof

1. We give a truth table for $p \Rightarrow p \vee q$ as follows.

p	q	$p \vee q$	$p \Rightarrow p \vee q$
1	1	1	**1**
1	0	1	**1**
0	1	1	**1**
0	0	0	**1**

Then $p \Rightarrow p \vee q$ is always true.

The truth table for $q \Rightarrow p \vee q$ is analogous to the one for $p \Rightarrow p \vee q$; the conclusion is the same.

2. In order to prove that $p \wedge q \Rightarrow p$ for all propositions p and q, we give a truth table for $p \wedge q \Rightarrow p$

p	q	$p \wedge q$	$p \wedge q \Rightarrow p$
1	1	1	**1**
1	0	0	**1**
0	1	0	**1**
0	0	0	**1**

Then $p \wedge q \Rightarrow p$ is always true.

The truth table for $p \wedge q \Rightarrow q$ is analogous to the one for $p \wedge q \Rightarrow p$.

1.3.4 Equivalence

Definition (Equivalence)

Two mathematical statements are **equivalent** if they have the same truth values.

The statement of the form "p if, and only if, q" is called an **equivalence** or **biconditional** statement. It is often abbreviated as p **iff** q and is written in symbols as $p \equiv q$ or $p \iff q$. It is equivalent to the compound statement "p implies q, and q implies p" composed of two CONDITIONAL statements. The truth-values of p and q must match for the biconditional statement as a whole to be true.

Examples:

1. "A triangle is equilateral if, and only if, it is equiangular" is a biconditional statement.

2. The proposition "$(1 = 1) \iff (0 = 0)$" is true, the proposition "$(1 = 0) \iff (2 = 0)$" is true, whereas the proposition "$(1 = 0) \iff (0 = 0)$" is false.

3. For all real x $(x \neq 0)$ and y , we have $y = x \iff \frac{y}{x} = 1$ is true.

4. The equivalence statement $(x = y \iff x^2 = y^2)$ is not true for all real x and y: for example $2^2 = (-2)^2$ $\not\Rightarrow 2 = -2$.

The truth-values of the basic connectives are given as follows:

p	q	$p \wedge q$	$p \vee q$	$p \Rightarrow q$	$p \Longleftrightarrow q$
1	1	1	1	1	1
1	0	0	1	0	0
0	1	0	1	1	0
0	0	0	0	1	1

Exercise 1: Prove the following equivalence by drawing the truth table:

$$p \Rightarrow q \Longleftrightarrow \overline{p} \vee q$$

Solution:

p	q	$\overline{p}$	$p \Rightarrow q$	$\overline{p} \vee q$
1	1	0	1	1
1	0	0	0	0
0	1	1	1	1
0	0	1	1	1

The truth table establishes that these corresponding pairs of compound statements are logically equivalent.

Definition (Contrapositive)

The contrapositive of $p \Rightarrow q$ is the proposition $\bar{q} \Rightarrow \bar{p}$. It can be shown that these two are equivalent:

$$(p \Rightarrow q) \Longleftrightarrow (\bar{q} \Rightarrow \bar{p})$$

The equivalence can easily be verified using truth table:

p	q	$\bar{p}$	$\bar{q}$	$p \Rightarrow q$	$\bar{q} \Rightarrow \bar{p}$
1	1	0	0	1	1
1	0	0	1	0	0
0	1	1	0	1	1
0	0	1	1	1	1

Logical Identities

★ **De Morgan's laws.**

1. $\overline{p \wedge q} \Longleftrightarrow \bar{p} \vee \bar{q}$

2. $\overline{p \vee q} \Longleftrightarrow \bar{p} \wedge \bar{q}$

Both of these laws can easily be verified using truth tables:

p	q	$\overline{p \wedge q}$	$\bar{p} \vee \bar{q}$
1	1	0	0
1	0	1	1
0	1	1	1
0	0	1	1

p	q	$\overline{p \vee q}$	$\bar{p} \wedge \bar{q}$
1	1	0	0
1	0	0	0
0	1	0	0
0	0	1	1

★ **Idempotence of $\wedge$ and $\vee$**

$P \Longleftrightarrow P \wedge P$ and $P \Longleftrightarrow P \vee P$

★ **Commutativity of $\wedge$ and $\vee$**

1. $p \wedge q \Longleftrightarrow q \wedge p$

2. $p \vee q \Longleftrightarrow q \vee p$

★ **Associativity of $\wedge$ and $\vee$**

1. $p \wedge (q \wedge r) \Longleftrightarrow (p \wedge q) \wedge r$

2. $p \vee (q \vee r) \Longleftrightarrow (p \vee q) \vee r$

★ **Distributivity of $\wedge$ over $\vee$ (and $\vee$ over $\wedge$ respectively)**

1. $p \wedge (q \vee r) \Longleftrightarrow (p \wedge q) \vee (p \wedge r)$

2. $p \vee (q \wedge r) \Longleftrightarrow (p \vee q) \wedge (p \vee r)$

★ **Domination laws**

1. $P \vee T \Longleftrightarrow T$

2. $P \wedge F \Longleftrightarrow F$

★ **Identity laws**

1. $P \vee F \Longleftrightarrow P$

2. $P \wedge T \Longleftrightarrow P$

★ **Negation laws**

1. $P \vee \bar{P} \Longleftrightarrow T$

2. $P \wedge \bar{P} \Longleftrightarrow F$

★ **Double negation law**

$$\bar{\bar{p}} \quad \Longleftrightarrow \quad p$$

★ **Absorption laws**

1. $P \vee (P \wedge Q) \Longleftrightarrow P$

2. $P \wedge (P \vee Q) \Longleftrightarrow P$

Exercise: Prove by applying the above rules.

a) $\overline{p \Rightarrow q} \Longleftrightarrow p \wedge \bar{q}$

b) $\bar{p} \Rightarrow \bar{q} \Longleftrightarrow p \vee \bar{q}$

Solution:

a) By applying De Morgan's laws:

$$\begin{aligned} \overline{p \Rightarrow q} \quad &\Longleftrightarrow \quad \overline{\bar{p} \vee q} \\ &\Longleftrightarrow \quad \bar{\bar{p}} \wedge \bar{q} \\ &\Longleftrightarrow \quad p \wedge \bar{q} \end{aligned}$$

b) By applying the following equivalence statement $p \Rightarrow q \Longleftrightarrow \bar{p} \vee q$, one has:

$$\begin{aligned} \bar{p} \Rightarrow \bar{q} \quad &\Longleftrightarrow \quad \bar{\bar{p}} \vee \bar{q} \\ &\Longleftrightarrow \quad p \vee \bar{q} \end{aligned}$$

Exercise: True or False. Prove by any method you like.

a) $p \Rightarrow (q \Rightarrow r) \Longleftrightarrow (p \Rightarrow q) \Rightarrow r$

b) $p \Rightarrow (q \vee r) \Longleftrightarrow (p \Rightarrow q) \vee (p \Rightarrow r)$

c) $p \wedge (q \Rightarrow r) \Longleftrightarrow (p \wedge q) \Rightarrow (p \wedge r)$

d) $p \vee (q \Rightarrow r) \Longleftrightarrow (p \vee q) \Rightarrow (p \vee r)$

Solution: For example, demonstrate the equivalence statement of (d) using a truth table (you will demonstrate the rest in a similar way)

p	q	r	$q \Rightarrow r$	$p \vee (q \Rightarrow r)$	$p \vee q$	$p \vee r$	$(p \vee q) \Rightarrow (p \vee r)$
1	1	1	1	**1**	1	1	**1**
1	1	0	0	**1**	1	1	**1**
1	0	1	1	**1**	1	1	**1**
1	0	0	1	**1**	1	1	**1**
0	1	1	1	**1**	1	1	**1**
0	1	0	0	**0**	1	0	**0**
0	0	1	1	**1**	0	1	**1**
0	0	0	1	**1**	0	0	**1**

We actually read the same truth values in the fifth and eighth columns.

You'll notice how we filled in the first three columns. This filling method makes it possible to forget no situation.

Exercise: If p and q are true and r and s are false statements, find the truth value of the following statements:

1. $(p \wedge q) \vee r$

2. $p \wedge (r \Rightarrow s)$

3. $(p \vee s) \Leftrightarrow (q \wedge r)$

4. $\overline{p \wedge \overline{r}} \vee (\overline{q} \vee s)$

Solution: Given that p and q are 1 and r and s are 0.

1. $(p \wedge q) \vee r \quad \Leftrightarrow \quad (1 \wedge 1) \vee 0$

$$\Leftrightarrow \quad 1 \vee 0$$

$$\Leftrightarrow \quad 1$$

♦ truth value of the given statement is true.

2. $p \wedge (r \Rightarrow s) \quad \Leftrightarrow \quad 1 \wedge (0 \Rightarrow 0)$

$$\Leftrightarrow \quad 1 \wedge 1$$

$$\Leftrightarrow \quad 1$$

♦ truth value of the given statement is true.

3. $(p \vee s) \Leftrightarrow (q \wedge r) \quad \Leftrightarrow \quad (1 \vee 0) \Leftrightarrow (1 \wedge 0)$

$$\Leftrightarrow \quad 1 \Leftrightarrow 0$$

$$\Leftrightarrow \quad 0$$

♦ truth value of the given statement is false.

4. $\overline{p \wedge \overline{r}} \vee (\overline{q} \vee s) \quad \Leftrightarrow \quad \overline{1 \wedge \overline{0}} \vee (\overline{1} \vee 0)$

$$\Leftrightarrow \quad 0 \vee 0$$

$$\Leftrightarrow \quad 0$$

♦ truth value of the given statement is false.

1.4 Predicates and Quantifiers

Definition (Predicate)

A **predicate** is a statement that contains variables and that may be true or false depending on the values of these variables.

Examples:

1. Let $P(x) : x^2 < x$ is a predicate. One has $P(1) : 1 < 1$ is false and $P(2) : 4 < 2$ is even false. But for $x = \frac{1}{2}$, $P(\frac{1}{2}) : \frac{1}{4} < \frac{1}{2}$ is true.

2. Let $P(x, y) : x^2 + y^2 = (x + y)^2$. Find the values of the following propositions: $P(0, 1)$, $P(0, 0)$, $P(1, 1)$. For which (x, y) is the value of $P(x, y)$ true?

A predicate can also be made a proposition by adding a **quantifier**. There are two quantifiers:

Definition (Universal quantifier)

A **universal quantifier** is a quantifier meaning "for all", "for any", "for each" or "for every", denoted by $\forall$.

Here is a formal way to say that for all values that a predicate variable x can take in a domain A, the predicate is true:

$$\underbrace{\forall\, x}_{} \quad \in A \quad , \quad P(x)$$

for all x belonging to A, $P(x)$ is true

Example: All natural numbers of the form $2n + 1$ are odd is written: $\forall\, n \in \mathbb{N}, 2n + 1$ is odd.

Definition (Existential quantifier)

An **existential quantifier** is a quantifier meaning "there exists", "there is at least one" or "for some".

Here is a formal way to say that for some values that a predicate variable x can take in a domain A, the predicate is true:

$$\underbrace{\exists\, x}_{} \quad \in A \quad , P(x)$$

for some x belonging to A, $P(x)$ is true

Example: There exists a natural number n satisfying $n \times n = n + n$ can be written: $\exists\, n \in \mathbb{N} : n \times n = n + n$.

Remark: A **unique existential quantifier** is a quantifier meaning "there is a unique", "there is exactly one" or "there exists only one". Here is a formal way to say that for some values that a predicate variable x can take in a domain A, the predicate is true:

$$\underbrace{\exists!\, x}_{} \quad \in A \quad , P(x)$$

there exists only one x belonging to A, $P(x)$ is true

Example: Let $P(x) : x + 2 = 5$.

1) $\forall x$, $P(x)$: "for all real numbers x, $x + 2 = 5$", which is false.

2) $\exists x$, $P(x)$: "there is a real number x such that $x + 2 = 5$", which is true.

3) $\exists! x$, $P(x)$: "there is a unique real number x such that $x + 2 = 5$", which is true.

Predicate Logic and Negating Quantifiers

We observe, at least intuitively, that the negations of $\exists$ and $\forall$ are correlated in the following manner.

$$\overline{\forall\, x, P(x)} \iff \exists\, x, \overline{P(x)}$$

$$\overline{\exists\, x, P(x)} \iff \forall\, x, \overline{P(x)}$$

Example: There is no natural number n satisfying $n \times n \times n = n + n + n$ as: $\overline{\exists\, n : n \times\ n \times\ n = n + n + n}$

Example: Let $P(x) : x + 2 = 5$.

$\overline{\exists x \in \mathbb{Z}, \text{“}x + 2 = 5\text{”}} \iff$ "$\forall x \in \mathbb{Z}, x + 2 \neq 5$"

Exercise: Write the negations by interchanging $\exists$ and $\forall$.

a) There is a real number x such that $x^2 < 0$.

b) Every integer is even.

c) There is an integer x such that $x^2 + 2x + 3 = 0$.

Solution:

a) $\overline{\exists x \in \mathbb{R}, \text{“}x^2 < 0\text{”}} \iff$ "$\forall x \in \mathbb{R}, x^2 \geq 0$"

b) There is an integer which is not even.

c) $\overline{\exists x \in \mathbb{Z}, \text{“}x^2 + 2x + 3 = 0\text{”}} \Longleftrightarrow \text{“}\forall x \in \mathbb{Z}, x^2 + 2x + 3 \neq 0\text{”}$

Exercise: Write the following proposition with quantifiers :

f is not increasing on $\mathbb{R}$ (where f is a function of $\mathbb{R}$ in $\mathbb{R}$).

Solution: By applying the negation of an implication studied above $[\overline{p \Rightarrow q} \Longleftrightarrow p \wedge \bar{q}]$, one has:

$$\overline{\forall\,(a,b) \in \mathbb{R}^2 / (a \leq b \Rightarrow f(a) \leq f(b))} \quad \Longleftrightarrow \quad \exists\,(a,b) \in \mathbb{R}^2, (a \leq b) \wedge (f(a) > f(b))$$

Exercise: Show that the function $\sin$ is not zero.

Solution: $\exists x = \frac{\pi}{2}, \sin(\frac{\pi}{2}) = 1 \neq 0$.Then $\sin \neq 0$.

1.4.1 Nested Quantifiers

Two quantifiers are nested if one is within the scope of the other. The order of **existential quantifiers** and **universal quantifiers** in a statement is important.

■ When we have one quantifier inside another, we need to be a little careful.

Example: Consider the following proposition over the integers:

$$\forall x \in \mathbb{Z}, \exists y \in \mathbb{Z} \;/\; (x + y = 0)$$

- The proposition is true.
- The existence of y depends on x : if you pick any x, I can find a y that makes $x + y = 0$ true.

Example: Consider the following proposition over the integers:

$$\exists y \in \mathbb{Z},\ \forall x \in \mathbb{Z} / \ (x + y = 0)$$

- The proposition is false.
- The existence of y does not depend on x : there is no y that will make $x + y = 0$ true for every x.

Example: Consider the following proposition over the integers:

$$\exists y \in \mathbb{Z},\ \forall x \in \mathbb{Z} / \ (x + y = x)$$

- The proposition is true.
- There is $y = 0$ that will make $x + y = x$ for every x.

Example: Suppose we claimed, "For every real number, there's a real number larger than it."

We'd write this as

$$\forall x \in \mathbb{R}, \exists y \in \mathbb{R} : y > x$$

- The proposition is true.
  - ■ We can exchange the same kind of quantifier ($\forall, \exists$).

These statements are equivalent:

$$\forall x, \forall y, P(x, y) \iff \forall y, \forall x, P(x, y)$$

$$\exists x, \exists y, P(x, y) \iff \exists y, \exists x, P(x, y)$$

Exercise: Translate the following statement into a logical expression. "Every real number except zero has a multiplicative inverse."

Solution:

$$\forall y \in \mathbb{R}^*, \exists x \in \mathbb{R} : xy = 1$$

Exercise: Express that the limit of a real-valued function f at point x_0 is l and express its negation: $\lim_{x \to x_0} f(x) = l$.

Solution:

In predicate logic:

$$\forall \varepsilon > 0, \exists\, \delta > 0, \forall x \in \mathbb{R} : (|x - x_0| < \delta \Rightarrow |f(x) - l| < \varepsilon)$$

its negation is given by: $\exists\, \varepsilon > 0, \forall \delta > 0, \exists\, x \in \mathbb{R} : |x - x_0| < \delta \wedge |f(x) - l| \geq 0$.

Exercise: If $A = \{3, 4, 6, 8\}$, determine the truth value of each of the following:

1. $\exists\, x \in A$, $x + 4 = 7$.

2. $\exists\, x \in A$, x is odd.

3. $\forall\, x \in A$, $(3 - x) \in \mathbb{N}$.

Solution:

1. Since $x = 3 \in A$, satisfies $x + 4 = 7$, the given statement is true. Its truth value is '1'.

2. Since $x = 3 \in A$, satisfies the given statement, the given statement is true. Its truth value is '1'.

3. $\exists\, x \in A, x = 4$, do not satisfy $3 - 4 = -1 \notin \mathbb{N}$,the given statement is false. Its truth value is '0'.

1.5 Methods of Proof

Our main interest in quantifiers for the purposes of this course is to develop techniques for proving mathematical statements.

When faced with a mathematical claim, understanding its quantifier is often a very good strategy for thinking about how to work out a proof.

Example: If the statement has the form $\forall x : P(x)$, then the global outline is likely to have the form: Consider any possible x, and show that it satisfies the property $P(x)$.

Example: If the statement has the form $\exists x : P(x)$, then the global outline is different: One needs to specify a particular x, and then show it satisfies $P(x)$.

1.5.1 Direct Method

We have already seen one way of proving a mathematical statement of the form: If p, then q. Based of the fact that the implication $p \Longrightarrow q$ is false only when p is true and q is false, the idea behind the method of proof that we discussed was to assume that p is true and then to proceed, through a chain of logical deductions, to conclude that q is true. Here is the outline of the argument:

Suppose that p is true.

$$\begin{aligned} p \quad &\Rightarrow r \\ &\Rightarrow s \\ &\Rightarrow .. \\ &\Rightarrow q \end{aligned}$$

Exercise: Prove the statement: If n is even, then n^2 is even.

Solution: Assume that the integer n is even.

$$\begin{aligned} \exists\, k \in \mathbb{Z}, n = 2k \quad &\Longrightarrow \qquad n^2 = (2k)^2 = 4k^2 \\ &\Longrightarrow \qquad n^2 = 2(2k^2) \\ &\Longrightarrow \quad n^2 = 2k^{'} \text{ such that } k^{'} = 2k^2 \end{aligned}$$

which shows that n^2 is even.

This is an example of a **direct method** of proof. In the following section we discuss **indirect methods** of proof.

1.5.2 Proof by Contrapositive

The idea behind this method of proof comes from the fact that the implication

$$\bar{q} \Rightarrow \bar{p}$$

is equivalent to the implication

$$p \Rightarrow q$$

Thus, in order to prove $p \Rightarrow q$, it suffices to prove: $\bar{q} \Rightarrow \bar{p}$. Here is the outline of the argument:

Suppose that : $\bar{q}$ is true

$$\begin{aligned} \bar{q} \quad &\Longrightarrow r \\ &\Longrightarrow s \\ &\Longrightarrow \ldots \\ &\Longrightarrow \bar{p}. \end{aligned}$$

Consequently, $\bar{q} \Rightarrow \bar{p}$ is true; therefore, $p \Rightarrow q$ is true.

Exercise: Prove that n^2 is even implies that n is even.

Solution: Suppose that n is not even. It then follows that

$$\begin{aligned} \exists\, k \in \mathbb{Z}, n = 2k+1 \quad &\Longrightarrow \quad n^2 = (2k+1)^2 = 4k^2 + 4k + 1 \\ &\Longrightarrow \quad n^2 = 2(2k^2 + 2k) + 1 \\ &\Longrightarrow \quad n^2 = 2k^{'} + 1 \text{ such that } k^{'} = 2k^2 + 2k \end{aligned}$$

Thus, n is not even implies that n^2 is not even, and therefore the contrapositive is true; namely, n^2 is even implies that n is even.

1.5.3 Proof by Contradiction (Absurd)

To prove that a proposition p is true we may assume that p is false then $\bar{p}$ is true. Therefore we show that it would lead to a **contradiction** or a false statement.

Exercise: Prove that $\sqrt{2}$ is irrational.

Solution: Let p: $\sqrt{2}$ is irrational. Now assume that p is false then $\bar{p}$ is true, that is, $\sqrt{2}$ is rational. Then there are some integers a and b with no common factors:

$$
\begin{aligned}
\exists\, a \in \mathbb{Z}, \exists\, b \in \mathbb{Z}^* : \sqrt{2} = a/b \quad &\Rightarrow \quad a^2 = 2b^2 \\
&\Rightarrow \quad a^2 \text{ is even} \\
&\Rightarrow \quad a = 2c, \;\; c \in \mathbb{Z} \\
&\Rightarrow \quad 4c^2 = 2b^2 \text{ (by Substituting)} \\
&\Rightarrow \quad 2c^2 = b^2 \\
&\Rightarrow \quad b^2 \text{ is even} \\
&\Rightarrow \quad b \text{ is even}
\end{aligned}
$$

This means that a and b have a common factor 2 which is a contradiction, and so $\bar{p}$ must be false and p is true.

1.5.4 Proof by Counter-Example

This proof structure allows us to prove that a property is not true by providing an example where it does not hold. Thus, in order to prove that the statement $\forall\, x,\, P(x)$ is false, it suffices to prove that the statement $\exists\, x, \overline{P(x)}$ is true.

Exercise: Prove that "all triangles are obtuse" is false.

Solution: We give the following counterexample: the equilateral triangle having all angles equal to sixty. In this case, there are infinitely many counterexamples. However, it only takes one.

Exercise: Prove that " If n is an integer and n^2 is divisible by 4, then n is divisible by 4" is false.

Solution: Consider $n = 6$. Then $n^2 = 36$ is divisible by 4, but $n = 6$ is not divisible by 4. Thus, $n = 6$ is a counterexample to the statement.

Exercise: Prove that "$(a + b)^2 = a^2 + b^2$" is not an algebraic identity, where $a, b \in \mathbb{R}$.

Solution: If $a = 1$ and $b = 2$, then $(a + b)^2 = 9$ and $a^2 + b^2 = 1^2 + 2^2 = 5$.

1.5.5 Proof by Cases Disjunction

This proof structure is used when one wants to prove a property $\forall\, x,\, P(x)$ depending on a parameter x belonging to a set A, and the proof depends on the value of x. Hence we decompose the set A into two or more sets $A_1, A_2, ...$ and we separate the reasonings following that $x \in A_1, x \in A_2, ...$ This proof is often used to solve (in) equations with absolute values (the proof depends on the sign of the quantity within the absolute value), to demonstrate properties in arithmetic (we separate the proof following the parity of some integers, their congruence modulo n ...).

To prove a proposition by case in the form $p \Rightarrow q$ where $p \Longleftrightarrow r \vee s$ we may instead prove both $r \Rightarrow q$ and $s \Rightarrow q$.

Exercise: Prove that for any integer n,the quotient $\frac{n(n+1)}{2}$ is an integer.

Solution:

- If n is even, then n is written $n = 2k$ and $n + 1 = (2k + 1)$. We then have $\frac{n(n+1)}{2} = k(2k + 1)$ which is an integer.
- If n is odd, then n is written $n = 2k + 1$ and $n + 1 = 2k + 2$. We then have $\frac{n(n+1)}{2} = (2k + 1)(k + 1)$ which is also an integer.

Exercise: Prove that $\forall x \in \mathbb{R} : |x - 1| \leq x^2 - x + 1$.

Solution: $|x-1| = \begin{cases} x-1 & \text{if } x \geq 1 \\ -x+1 & \text{if } x < 1 \end{cases}$

$$|x-1| \leq x^2 - x + 1 \Leftrightarrow \begin{cases} x - 1 \leq x^2 - x + 1 & \text{if } x \geq 1 \\ -x + 1 \leq x^2 - x + 1 & \text{if } x < 1 \end{cases}$$

1. If $x \geq 1, x - 1 - x^2 + x - 1 \leq 0 \Leftrightarrow -x^2 + 2x - 2 \leq 0$ is true because the discriminant of the equation $x^2 - 2x + 2 = 0$ is negative ($\triangle = -4$), hence $x^2 - 2x + 2 \geq 0$

2. If $x < 1, -x + 1 - x^2 + x - 1 \leq 0 \Leftrightarrow -x^2 \leq 0$ is true.

Therefore $\forall x \in \mathbb{R} : |x - 1| \leq x^2 - x + 1$.

1.5.6 Proof by Mathematical Induction

To prove a proposition in the form $\forall n \in \mathbb{N},\ n \geq n_0$, $P(n)$ where n is a natural number, it suffices to prove it in two steps:

1. $P(n_0)$ is true for a certain base step n_0. Usually, the base case is $n = 1$ or $n = 0$.

2. $P(n) \Rightarrow P(n+1)$. That is, if $P(n)$ is true, then $P(n+1)$ is true.

Exercise: Prove the following formula for all natural numbers n.

$1 + 3 + 5 + 7 + 9 + ... + (2n - 1) = n^2$.

Solution: Let $P(n) : 1 + 3 + 5 + 7 + 9 + ... + (2n - 1) = n^2$

We shall prove $\forall n \in \mathbb{N}, P(n)$ in two steps:

1) $P(0) : 0 = 0^2$ so this proposition is true.

2) Let $P(n) : 1 + 3 + 5 + 7 + 9 + ... + (2n - 1) = n^2$

$$\Rightarrow 1 + 3 + 5 + 7 + 9 + ... + (2n - 1) + (2n + 1) = n^2 + (2n + 1)$$

$$\Rightarrow 1 + 3 + 5 + 7 + 9 + ... + (2n - 1) + (2n + 1) = (n + 1)^2$$

$$\Rightarrow 1 + 3 + 5 + 7 + 9 + ... + (2n - 1) + (2(n + 1) - 1) = (n + 1)^2$$

$$\Rightarrow P(n + 1) \text{ is true}$$

Therefore $\forall n \in \mathbb{N}, P(n)$

Exercise: is $3^n - 1$ a multiple of 2 ?

Solution:

1. Show it is true for $n = 1$, $3^1 - 1 = 3 - 1 = 2$. One has 2 is a multiple of 2. That was easy. $3^1 - 1$ is a multiple of 2.

2. Assume it is true for n and prove that $3^{n+1} - 1$ is a multiple of 2?;

$$
\begin{aligned}
3^n - 1 = 2k \quad &\Rightarrow \quad 3^n \times 3 - 1 \times 3 = 2k \times 3 \\
&\Rightarrow \quad 3^n \times 3 - 3 = 2k \times 3 \\
&\Rightarrow \quad 3^{n+1} - 1 = 2 + 2k \times 3 \\
&\Rightarrow \quad 3^{n+1} - 1 = 2(1 + 3k) \\
&\Rightarrow \quad 3^{n+1} - 1 = 2k^{'} \text{ such that } k^{'} = 1 + 3k
\end{aligned}
$$

Therefore $\forall n \in \mathbb{N}^*,\ 3^n - 1$ a multiple of 2

FAMOUS ISLAMIC MATHEMATICIANS



Omar Khayyam

Omar Khayyam wrote one of the most famous treatises of **Algebra**. He created the first theories known about parallels in quadrilaterals. He also found out how to solve roots of any degree, laying down some of the principles of Algebra . He had notable works in **geometry**, particularly on the theory of proportions

Chapter 2

Sets, Relations and Applications

2.1 Set Theory

Set is a one of the most basic concept in mathematics, since we almost always have a collection of things we are trying to study. In particular, mathematical structure arises from operations and relationships, such as addition, multiplication and in the case of an application. Set is used in all branches of mathematics and computer science. In analysis branch for example, an understanding of limit points and what is meant by the continuity of a function, differential and integral calculus are based on set theory.

2.1.1 Relationships between elements and parts of a set

Definition (Set, Element)

A **set** A is a collection of objects called the **elements** of the set.

- If x is an element of the set A then we write $x \in A$, while the negation is written $x \notin A$.
- Set is typically specified either *explicitly*, that is by listing all the elements the set contains, or *implicitly*, using a predicate description as seen in predicate logic, of the form $\{x|\ P(x)\}$.
- The ordering of the elements is not important and repetition of elements is ignored.
- A set may also be **empty** set (or **null set**) and it is denoted by $\emptyset$ (phi) or $\{\}$.
- The **universe** E is the biggest set in which all the other sets we are interested in lie.

Examples:

1. The set A given by $\{1, 2, 3\}$ is an explicit description.

2. The set $\{x,\ x$ is a prime number $\}$ is implicit.

3. $\{\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}\}$ is a set containing five sets.

4. $\{x : x \in \{2, 3, 5\}$ and $x \leq 1\}$ is an empty set.

5. $\{x : x^2 = -1\ \}$ is the set of two elements: i and $-i$.

6. $\{e, \pi, 1, \pi, 2, 1\} = \{1, 2, \pi, e\}$

Definition (Cardinality)

If a set A contains exactly n elements where n is a non-negative integer, then A is a finite set, and n is called the **cardinality** of A. We write $|A| = n$.

Remark: If $|A|$ is finite, A is a finite set; otherwise, A is infinite.

Examples:

1. $A = \{1, 2, \sqrt{7}, 0\}, |A| = 4$.

2. $|\{x \mid\ -2 < x < 5, x \in \mathbb{Z}\}| = 6$.

3. $|\emptyset| = 0$.

4. $|\{x \mid (x \in \emptyset) \wedge (x < -4)\}| = 0$.

5. The set of positive integers is an infinite set.

Using set notation with quantifiers

Sometimes, we restrict the domain of a quantified statement explicitly by using set notations.

- We use $\forall\, x \in A$, $(P(x))$ to denote that $P(x)$ **holds for every** $x \in A$.

- We use $\exists\, x \in A$, $(P(x))$ to denote that $P(x)$ **holds for some** $x \in A$.

Definition (Equality)

Two sets A and B are **equal** if each element of A is an element of B and vice versa. This is denoted, $A = B$. Formally:

$$\boxed{A = B \Leftrightarrow \forall x : x \in A \Leftrightarrow x \in B} \tag{2.1}$$

Examples:

1. $\{1, 2, 3\} = \{2, 1, 3\}$.
2. $\{1, 2, 3, 4\} = \{x \in \mathbb{N}\,, x < 5\}$.
3. $\{x \in \mathbb{R} : x^2 + 1 = 0\} = \emptyset$.

To say that two sets A and B are *not equal*, inequality is $A \neq B$ of course. We use the negation from predicate logic, which is (using the rules we have studied in predicate logic! namely negation of universal quantifier and De Morgan's law). One obtains:

$$\begin{aligned}
\overline{\forall x : x \in A \Leftrightarrow x \in B} \quad &\Leftrightarrow \quad \exists\, x : \overline{(x \in A \Rightarrow x \in B) \wedge (x \in B \Rightarrow x \in A)} \\
&\Leftrightarrow \quad \exists\, x : \overline{(x \in A \Rightarrow x \in B)} \vee \overline{(x \in B \Rightarrow x \in A)} \\
&\Leftrightarrow \quad \exists\, x : \left(\overline{\overline{(x \in A)} \vee (x \in B)}\right) \vee \left(\overline{\overline{(x \in B)} \vee (x \in A)}\right) \\
&\Leftrightarrow \quad \exists\, x : \left((x \in A) \wedge \overline{(x \in B)}\right) \vee \left((x \in B) \wedge \overline{(x \in A)}\right) \\
&\Leftrightarrow \quad \exists\, x : ((x \in A) \wedge (x \notin B)) \vee ((x \in B) \wedge (x \notin A))
\end{aligned}$$

Definition (Subset)

- A set A is a **subset** of B *if and only if* every element of A is also in B. We use $A \subseteq B$ to indicate A is a subset of B , that means A is **included** in B. Formally

$$\boxed{A \subseteq B \Leftrightarrow \forall x : x \in A \Rightarrow x \in B} \tag{2.2}$$

- A is a **proper subset (or strict subset)** of B, $A \subset B$, if $A \subseteq B$ and $A \neq B$.

- Note the difference between $x \in A$ and $\{x\} \subseteq A$: in the first expression, x is an *element* of A, while in the second, we consider the *subset* $\{x\}$, which is emphasized by the bracket notation.

- To say that A is *not a subset* of B, we use the negation of the following statement $[\forall x : x \in A \Rightarrow x \in B]$, which is:

$$\overline{\forall x : x \in A \Rightarrow x \in B} \quad \Leftrightarrow \quad \exists\, x : (x \in A) \wedge (x \notin B)$$

Therfore,

$$\boxed{A \nsubseteq B \Leftrightarrow \exists\, x : (x \in A) \wedge (x \notin B)} \tag{2.3}$$

Examples:

1. $\{12, 43, 66\} \subseteq \{12, 43, 66\}$

2. $\{a, \bigstar\} \subset \{a, b, \bigstar, \blacktriangle\}$

3. $\mathbb{N} \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R}$

4. Let $A = \left\{\sqrt{2}, i\right\}$, $A \nsubseteq \mathbb{R}$; $i \in A$ and $i \notin \mathbb{R}$

Remark: There is a difference between $\emptyset$ and $\{\emptyset\}$: the first one is an empty set, the second one is a set, which is not empty since it contains one element: the empty set!

Properties: Notice that $A \subseteq A$ and in fact each set is a subset of itself. The empty set $\emptyset$ is a subset of any set $\emptyset \subseteq A$.

Exercise: Prove that $\emptyset \subseteq A$

Solution: Recall the definition of a subset: all elements of a set A must be also elements of B; $\forall x : x \in A \Rightarrow x \in B$. We must show the following implication is true for any $A, \forall x : x \in \emptyset \Rightarrow x \in A$. Since the empty set does not contain any element, $x \in \emptyset$ is *always False statement*. Then the implication is *always True*.

Venn Diagram

A diagram in which mathematical sets are represented by overlapping circles within a boundary representing the **universal set** is called a **Venn diagram**. Such diagrams provide convenient pictorial representations of

relations between sets.

Example: In the diagram (See Figure 2.1) a **universal set** E is represented by the interior of a rectangle, and one subset A of E as the interior of one overlapping circle within the rectangle.



Figure 2.1

The Venn diagram in Figure 2.1 shows: $A \subseteq E$, $\{1, 2, 3, \pi\} \subseteq E$, $\{1, 2, 3, \pi\} \nsubseteq A$, ...

Definition (Power set)

Given a set E, the power set of E is the set of all subsets of E. The power set is denoted by $P(E)$. Formally:

$$\boxed{P(E) = \{A,\ A \subseteq E\ \}} \tag{2.4}$$

Examples: Write the power set of the following sets: $\emptyset$, $\{1\}$, $\{1, 2\}$, $\{1, 2, 3\}$. If E is a set with $|E| = n$ then $|P(E)| = ?$

1. $P(\emptyset) = \{\emptyset\}$ and $|P(\emptyset)| = 1$.

2. $P(\{1\}) = \{\emptyset, \{1\}\}$ and $|P(\{1\})| = 2$.

3. $P(\{1, 2\}) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$ and $|P(\{1, 2\})| = 4$.

4. $P(\{1, 2, 3\}) = \{\emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$ and $|P(\{1, 2, 3\})| = 8$.

Property: If E is a set with $|E| = n$ then $|P(E)| = 2^n$.

2.1.2 Set Operations

There are a number of basic set manipulations, each of which can be depicted with a VENN DIAGRAM.

Definition (Set Intersection)

The **intersection** of the sets A and B is the set of all elements that are in both A and B. We write $A \cap B$.

Formally:

$$\boxed{A \cap B \quad = \quad \{x, (\, x \in A) \;\; \wedge \;\; (x \in B)\}} \tag{2.5}$$

Venn Diagram of Intersection Operation (See figure 2.2):



Figure 2.2

Example:

1. $\{1, 2, 3, 4\} \cap \{-3, 4, 5\} = \{4\}$

2. $\{x|\; x > 0\} \cap \{x|\; x \geq 2\} = \{x|\; x \geq 2\}$.

3. $\mathbb{N} \cap \mathbb{Z} \cap \mathbb{R} = \mathbb{N}$

Definition (Set Disjoint)

Two sets A and B are disjoint if $A \cap B = \emptyset$.

Examples:

1. $\{2, 4, 6\} \cap \{8, 10, 12\} = \emptyset$, so they are disjoint.

2. $\{1, 2, 3\} \cap \{3, 4, 5\} \neq \emptyset$, so they are not disjoint.

3. $\mathbb{N} \cap \mathbb{Z} \neq \emptyset$, so they are not disjoint.

4. $\{x|\ x \geq 1\} \cap \mathbb{Z}^- = \emptyset$, so they are disjoint.

Definition (Set Union)

The **union** of two sets A and B , denoted by $A \cup B$, is the set that contains exactly all the elements that are in either A or B (or in both). Formally:

$$\boxed{A \cup B \quad = \quad \{x,\ (x \in A)\ \vee\ (x \in B)\}} \tag{2.6}$$

Venn Diagram of Union Operation (See Figure 2.3):



Figure 2.3

Examples:

1. Let $A = \{0, 1, 2, 3, 6\}$, $B = \{0, 1, 2, 4, 6, 9\}$, $A \cup B = \{0, 1, 2, 3, 4, 6, 9\}$.

2. $\mathbb{Z}^- \cup \mathbb{Z}^+ = \mathbb{Z}$.

3. $\{x|\ x > 0\} \cup \{x|\ x > -1\} = \{x|\ x > -1\}$.

Lemma (Cardinality of intersection and union)

For any two sets A and B, we have

$$\boxed{|A \cup B| \quad = \quad |A| + |B| - |A \cap B|} \tag{2.7}$$

Theorem 2. If A and B are any sets, then

1. $(A \cap B) \subseteq A$ and $(A \cap B) \subseteq B$

2. $A \subseteq (A \cup B)$ and $B \subseteq (A \cup B)$

Proof

1) To prove that $A \cap B \subseteq A$, we must show that $x \in A \cap B \Rightarrow x \in A$.

$$\begin{aligned} x \in A \cap B \quad &\Rightarrow \quad (x \in A) \wedge (x \in B) && \text{by (2.5)} \\ &\Rightarrow \quad x \in A && \text{by theorem 1} \end{aligned}$$

Analogously, we can show that $(A \cap B) \subseteq B$.

2) To prove that $A \subseteq A \cup B$, we must show that $x \in A \Rightarrow x \in A \cup B$:

$$\begin{aligned} x \in A \quad &\Rightarrow \quad (x \in A) \vee (x \in B) && \text{by theorem 1} \\ &\Rightarrow \quad x \in A \cup B && \text{by (2.6)} \end{aligned}$$

Analogously, we can show that $B \subseteq A \cup B$

Properties: For all subsets A, B and C of the univers E , the following are true.

- **Commutative laws:** $\begin{cases} A \cap B = B \cap A \\ A \cup B = B \cup A \end{cases}$

- **Associative laws:** $\begin{cases} (A \cap B) \cap C = A \cap (B \cap C) \\ (A \cup B) \cup C = A \cup (B \cup C)\,. \end{cases}$

- **Distributive laws:** $\begin{cases} A \cap (B \cup C) &= (A \cap B) \cup (A \cap C) \\ A \cup (B \cap C) &= (A \cup B) \cap (A \cup C) \end{cases}$

- **Identity laws:** $\begin{cases} A \cup \emptyset &= A \\ A \cap E &= A \end{cases}$

- **Domination laws:** $\begin{cases} A \cup E &= E \\ A \cap \emptyset &= \emptyset \end{cases}$

- **Idempotent laws:** $\begin{cases} A \cup A &= A \\ A \cap \emptyset &= \emptyset \end{cases}$

Definition (Set Partition)

A collection of nonempty sets $\{A_1,\ A_2, ...,\ A_n\}$ is a partition of a set A if and only if

1. $A = A_1 \cup\ A_2 \cup ...\ \cup\ A_n$.

2. $A_1,\ A_2, \cdots,\ A_n$ are mutually disjoint (or pairwise disjoint) : $Ai \cap Aj = \emptyset, i \neq j, i, j = 1, 2, .., n$.

Example: Consider $A = \mathbb{Z}$, $A_1 = \{x,\ x \text{ is even}\}$, $A_2 = \{x,\ x \text{ is odd}\}$.

Then A_1, A_2 form a partition of A.

Venn Diagram of Set partition (See Figure 2.4):

Definition (Set difference)

The **difference** of A and B, is the set containing elements that are in A but not in B. Formally:

$$\boxed{A - B \quad = \quad \{x,\ (x \in A) \wedge (x \notin B)\}} \tag{2.8}$$

Venn Diagram of Set difference (See Figure 2.5):



Figure 2.4



Figure 2.5

Examples:

1. $\{1,2,3\} - \{3,4,5\} = \{1,2\}$.

2. $\mathbb{R} - \{0\} = \{x | \ (x \in \mathbb{R}) \wedge (x \neq 0)\}$.

3. $\mathbb{N} - \left\{\frac{a}{b}, (a \in \mathbb{Z}) \wedge (b \in \mathbb{Z}^*)\right\} = \mathbb{N}$.

Properties: Let A and B subsets of the univers E.

- $A - B \subset A$
- $A - A = \emptyset, A - \emptyset = A, \emptyset - A = \emptyset$
- $A - (A - B) = A \cap B$
- $A - B = \Phi \Leftrightarrow A \subseteq B$

Definition (Set complement)

Let A subset of the universal set E. The complement of set A with respect to E, denoted by C_E^A or CA or $\overline{A}$, is the set that contains exactly all the elements that are not in A. Formally:

$$\boxed{\overline{A} \;=\; E - A \;=\; \{x \in E / \; x \notin A\}} \tag{2.9}$$

Venn Diagram of Set complement (See Figure 2.6):



Figure 2.6

Examples: Let the universe be $\mathbb{R}$

1. $\overline{\{0\}} = \{x,\ x \neq 0\} = \mathbb{R}^*$.

2. $\overline{\mathbb{R}^-} = \{x,\ x > 0\} = \mathbb{R}^+$.

3. $\overline{]-1, 2]} =]-\infty, -1] \cup]2, +\infty[$.

4. $\overline{\mathbb{R}} = \{x,\ x \notin \mathbb{R}\} = \emptyset$.

Properties: Let A and B subsets of the univers E.

- $\overline{E} = C_E^E = \emptyset, \overline{\emptyset} = C_E^{\emptyset} = E$
- $C_E\left(C_E^A\right) = \overline{\overline{A}} = A$
- $A \cup \overline{A} = E, A \cap \overline{A} = \emptyset$
- $A \subset B \Longrightarrow \overline{B} \subset \overline{A}$

DE MORGAN'S LAWS

Explain how set complement interacts with intersections and unions of sets.

$$\begin{cases} \overline{A \cap B} &= \overline{A} \cup \overline{B} \\ \\ \overline{A \cup B} &= \overline{A} \cap \overline{B} \end{cases}$$

Exercise: Let A and B are subsets of the universal set E. Show that:

1. $C_E\left(C_E^A\right) = \overline{\overline{A}} = A$
2. $A \subset B \Longrightarrow \overline{B} \subset \overline{A}$
3. $\overline{A \cap B} = \overline{B} \cup \overline{A}$
4. $\overline{A \cup B} = \overline{B} \cap \overline{A}$

Solution:

1. $C_E\left(C_E^A\right) = \overline{\overline{A}} = A$?

   $x \in C_E\left(C_E^A\right) \Longleftrightarrow x \notin \left(C_E^A\right) \Longleftrightarrow x \in A$

2. $A \subset B \Longrightarrow \overline{B} \subset \overline{A}$?

$$A \subset B \quad \Longleftrightarrow (\forall x, x \in A \Longrightarrow x \in B)$$

$$\Longleftrightarrow (\forall x, x \notin B \Longrightarrow x \notin A)$$

$$\left(\forall x, x \in \overline{B} \Longrightarrow x \in \overline{A}\right)$$

$$\overline{B} \subset \overline{A}$$

3. $\overline{A \cap B} = \overline{B} \cup \overline{A}$

$$x \in \overline{A \cap B} \qquad \Longleftrightarrow x \notin (A \cap B)$$

$$\Longleftrightarrow (x \notin A) \vee (x \notin B)$$

$$\Longleftrightarrow \big(x \in \overline{A}\big) \vee \big(x \in \overline{B}\big)$$

$$\Longleftrightarrow x \in \overline{A} \cup \overline{B}$$

4. $\overline{A \cup B} = \overline{B} \cap \overline{A}$

$$x \in \overline{A \cup B} \qquad \Longleftrightarrow x \notin (A \cup B)$$

$$\Longleftrightarrow (x \notin A) \wedge (x \notin B)$$

$$\Longleftrightarrow \big(x \in \overline{A}\big) \wedge \big(x \in \overline{B}\big)$$

$$\Longleftrightarrow x \in \overline{A} \cap \overline{B}$$

Definition (Set Symmetric Difference)

The **symmetric difference** of set A and set B, denoted by $A \triangle B$, is the set containing those elements in exactly one of A and B.

Formally:

$$\boxed{A \triangle B \quad = \quad (A - B) \cup (B - A)} \tag{2.10}$$

Venn Diagram of Set difference (See Figure 2.7):



Figure 2.7

Properties: Let A and B subsets of the univers E.

- $A \triangle B = B \triangle A$
- $A \triangle \emptyset = A$
- $A \triangle E = E - A$
- $A \triangle A = \emptyset$
- $A \triangle B = (A \cup B) - (A \cap B)$
- $A \triangle B = \overline{A} \triangle \overline{B}$
- $(A \triangle B) \triangle C = A \triangle (B \triangle C)$

Example: If $A = \{1, 2, 3, 4, 5, 10\}$ and $B = \{0, 1, 2, 3, 4, 5, 7, 8, 9\}$, then $A - B = \{10\}$ and $B - A = \{0, 7, 8, 9\}$. Hence $A \triangle B = \{0, 7, 8, 9, 10\}$

Definition (Ordered tuple)

An **ordered** n**-tuple** $(x_1, x_2, ..., x_n)$ has x_1 as its first element, x_2 as its second element, . . ., x_n as its nth element. The order of elements is important in such a tuple. Note that $(x_1, x_2) \neq (x_2, x_1)$ but $\{x_1, x_2\} = \{x_2, x_1\}$.

Definition (Set Cartesian product)

The **Cartesian product** of the sets A and B, denoted by $A \times B$ is the set of all ordered pairs (x_1, x_2), where $x_1 \in A$, $x_2 \in B$:

$$\boxed{A \times B = \{(x_1, x_2)/ \quad x_1 \in A, x_2 \in B \ \}} \tag{2.11}$$

The equality in $A \times B$ is defined by: $(x_1, y_1) = (x_2, y_2) \Longleftrightarrow x_1 = x_2 \wedge y_1 = y_2$.

Cartesian product can be formed from n sets $A_1, A_2, ..., A_n$, denoted by $A_1 \times A_2 \times \cdots \times A_n$, is defined as the set of ordered tuples $(x_1, x_2, ..., x_n)$ where $x_1 \in A_1, x_2 \in A_2, ..., x_n \in A_n$. That is:

$$\boxed{A_1 \times A_2 \times \cdots \times A_n = \{(x_1, x_2, .., x_n)/ \ x_1 \in A_1, x_2 \in A_2, .., x_n \in A_n\}} \tag{2.12}$$

$$\boxed{A_1 = A_2 = \cdots = A_n = A \quad \Longrightarrow \quad A_1 \times A_2 \times \cdots \times A_n = A^n} \tag{2.12}$$

If we represent a set $A \times B$, then a segment of the horizontal axis is marked off to represent A and a segment of the vertical axis is marked off to represent B; $A \times B$ is the rectangle determined by these two segments (See Figure 2.8).



Figure 2.8

Examples: Let $A = \{-2, 3\}$ and $B = \{0, -4, 2\}$

1. $A \times B = \{(-2, 0), (-2, -4), (-2, 2), (3, 0), (3, -4), (3, 2)\}$.
2. $B \times A = \{(0, -2), (0, 3), (-4, -2), (-4, 3), (2, -2), (2, 3)\}$
3. Note that $A \times B \neq B \times A$.
4. $\mathbb{R} \times \mathbb{R} = \{(x, y) \,|x \in \mathbb{R}, y \in \mathbb{R}\}$ is the set of point coordinates in the 2D Plane.
5. $\mathbb{R} \times \mathbb{R} \times \mathbb{R} = \{(x, y, z) \,|x \in \mathbb{R}, y \in \mathbb{R}, z \in \mathbb{R}\}$ is the set of point coordinates in the 3D Space.

Lemma (**Cardinality of Cartesian product**)

In general, if A_i's are finite sets, we have:

$$\boxed{|A_1 \times \ A_2 \times ... \times \ A_n| = |A_1| \times \ |A_2| \times ... \times \ |A_n|} \tag{2.13}$$

Properties: If A and B are any sets, then

- $A \times B \neq B \times A$
- $A \times \emptyset = \emptyset \times A = \emptyset$
- $A \subseteq E \wedge B \subseteq F \Leftrightarrow A \times B \subseteq E \times F$
- $A \times (B \cap C) = (A \times B) \cap (A \times C)$
- $A \times (B \cup C) = (A \times B) \cup (A \times C)$
- $A \times B = \emptyset \Leftrightarrow A = \emptyset \vee B = \emptyset$
- $A \times B \neq \emptyset \Leftrightarrow A \neq \emptyset \wedge B \neq \emptyset$

Exercise : The sets $A = \{1, 2, x\}$, $B = \{3, 4, y\}$ are given. Determine x and y, knowing that $\{1, 3\} \times \{2, 4\} \subseteq A \times B$

Solution: We form the sets $A \times B$ and $C = \{1, 3\} \times \{2, 4\}$:

$A \times B = \{(1, 3), (1, 4), (1, y), (2, 3), (2, 4), (2, y), (x, 3), (x, 4), (x, y)\}$

$C = \{(1, 2), (1, 4), (3, 2), (3, 4)\}$

Because $\{1, 3\} \times \{2, 4\} \subseteq A \times B$, we obtain

$(1, 2) \in C \Longrightarrow (1, 2) \in A \times B \Longrightarrow (1, 2) = (1, y) \Longrightarrow y = 2.$

$(3, 4) \in C \Longrightarrow (3, 4) \in A \times B \Longrightarrow (3, 4) = (x, 4) \Longrightarrow x = 3.$

For $x = 3$ and $y = 2$, we have $(3, 2) \in A \times B$.

Therefore: $x = 3$ and $y = 2$.

Exercise : Determine the sets A and B that simultaneously satisfy the following conditions:

1. $A \cup B = \{1, 2, 3, 4, 5\}$;
2. $A \cap B = \{3, 4, 5\}$;
3. $2 \notin (B - A)$

4. $1 \notin (A - B)$

Solution:

$1 \notin (A - B) \Leftrightarrow (1 \notin A) \vee (1 \in B)$

$1 \in (A \cup B) \Leftrightarrow (1 \in A) \vee (1 \in B)$

$$\begin{aligned}
[1 \notin (A - B)] \wedge [1 \in (A \cup B)] &\Leftrightarrow [(1 \notin A) \vee (1 \in B)] \wedge [(1 \in A) \vee (1 \in B)] \\
&\Leftrightarrow [(1 \notin A) \wedge (1 \in A)] \vee (1 \in B) \\
&\Leftrightarrow F \vee (1 \in B) \\
&\Leftrightarrow 1 \in B
\end{aligned}$$

$2 \notin (B - A) \Leftrightarrow (2 \notin B) \vee (2 \in A)$

$2 \in (A \cup B) \Leftrightarrow (2 \in A) \vee (2 \in B)$

$$\begin{aligned}
[2 \notin (B - A)] \wedge [2 \in (A \cup B)] &\Leftrightarrow [(2 \notin B) \vee (2 \in A)] \wedge [(2 \in A) \vee (2 \in B)] \\
&\Leftrightarrow [(2 \notin B) \wedge (2 \in B)] \vee (2 \in A) \\
&\Leftrightarrow F \vee (2 \in A) \\
&\Leftrightarrow 2 \in A
\end{aligned}$$

Then, $A = \{2, 3, 4, 5\}$ and $B = \{1, 3, 4, 5\}$

2.2 Relations

The notion of relation is omnipresent, in mathematics as in everyday life. The intuitive idea is to understand the fact that a certain link exists or not between two or more objects.

The concept of relation finds a precise characterization in a mathematical context, the Cartesian product operation offering in this respect a frame both propitious and fertile.

2.2.1 Binary Relations

Binary relations are an excellent way for capturing certain structures that appear in computer science.

Definition (Binary Relation)

A **binary relation** over a nonempty set A is a predicate R that can be applied to ordered pairs (x, y) of elements x and y given from A.

The representing **graph** of a relation in A is a graph $G \subseteq A \times A$ which consists of all the pairs (x, y) such that the relation between two elements x and y is true. Conversely, if we are given an arbitrary graph $G \subseteq A \times A$, then G defines a relation in A, namely the relation R is true if and only if $(x, y) \in G$.

Notation for Binary Relations

Let R be a binary relation in A . Then

$$\boxed{x\ R\ y \Longleftrightarrow (x, y) \in G}$$

Examples:

1. "x is greater than y ".
2. "x and y have the same absolute value".
3. "$x^2 + y^2 = 1$".
4. $A \subset B$.

Example: Suppose $A = \{1, 2, 3, 4\}$. We give the graph $G \subseteq A \times A$ of the following relation:

$\forall\, x,\ y \in A: \quad x\ R\ y \quad \Longleftrightarrow \quad x < y.$

Then, $G = \{(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4)\}$.

Example: A relation R is defined on $\mathbb{R}$ by:

$\forall\, x,\ y \in \mathbb{R}: \quad x\ R\ y \quad \Longleftrightarrow \quad xy^3 - x^3y = 6$

We show that $1\ R\ 2$, because $1 \times 2^3 - 1^3 \times 2 = 6$

Remark: If R is a binary relation over A and it does not hold for the pair (x, y), then $\overline{x\ R\ y}$.

Examples: $3 \neq 4, \mathbb{R} \nsubseteq \mathbb{Z}, 4 \nleqslant 3$.

Properties of a relation

Let R be a binary relation in A.

- R is **reflexive** iff $\forall x \in A: \ x \, R \, x$.
- R is **symmetric** iff $\forall x, y \in A : x \, R \, y \Rightarrow y \, R \, x$.
- R is **anti-symmetric** iff $\forall x, y \in A : (x \, R \, y) \wedge (y \, R \, x) \Longrightarrow x = y$.
- R is **transitive** iff $\forall x, y, z \in A : (x \, R \, y) \wedge (y \, R \, z) \Longrightarrow x \, R \, z$.

Example: Let $A = \{1, 2, 3\}$ and consider three relations R, T, S on A :

$G_R = \{(1, 1), (1, 2), (2, 1), (2, 2), (3, 3)\}$.

$G_S = \{(1, 1), (1, 3), (2, 2), (3, 2)\}$.

$G_T = \{(1, 2), (1, 3), (2, 3)\}$.

- R is reflexive, symmetric, and transitive, but not anti-symmetric because $(1 \, R \, 2) \wedge (2 \, R \, 1)$ but $1 \neq 2$.
- S is anti-symmetric, but not reflexive because $\left(\overline{3 \, R \, 3}\right)$, not symmetric $(1 \, R \, 3 \,)$ but $\overline{(3 \, R \, 1)}$, and not transitive $(1 \, R \, 3) \wedge (3 \, R \, 2)$ but $\left(\overline{1 \, R \, 2}\right)$.
- T is anti-symmetric and transitive, but not reflexive $\left(\overline{1 \, R \, 1}\right)$ and not symmetric $(1 \, R \, 2 \,)$ but $\overline{(2 \, R \, 1)}$.

2.2.2 Equivalence Relation

Definition (Equivalence Relation)

An **equivalence relation** is a relation that is *reflexive, symmetric and transitive.*

Examples:

1. The "*equal-to*" relation, " $=$ ", on $\mathbb{R}$ is an equivalence relation.
2. The "*less- than -or- equal to* " relation, "$\leqslant$", on $\mathbb{R}$ is not an equivalence relation because it is not symmetric. For example: $1 \leqslant 2$ but $2 \nleqslant 1$.
3. The "*strictly-less-than*" relation, "$<$ ", on $\mathbb{R}$ is not an equivalence relation because it is not reflexive. For example: $1 \nless 1$.

4. The "*Line Parallel relation*", "$\|$" , is an equivalence relation.

5. The "*Perpendicular Lines"* relation, "$\perp$", is symmetric but neither reflexive nor transitive.

6. The "subset" relation "$\subseteq$ ", on $P(E)$ such that $E = \{1, 2, 3\}$, is not an equivalence relation because it is not symmetric. For example: $\{1\} \subseteq \{1, 2\}$ but $\{1, 2\} \nsubseteq \{1\}$.

Exercise: Let A be the set of all triangles in a plane with R a relation in A given by

$$\forall\, T_1,\ T_2 \ \in A: \quad T_1\ R\, T_2 \quad \Longleftrightarrow \quad T_1 \text{ is congruent to } T_2.$$

Show that R is an equivalence relation.

Solution:

1. R is reflexive, since every triangle is congruent to itself.

2. $T_1\ R\, T_2 \Longrightarrow T_1$ is congruent to $T_2 \Longrightarrow T_2$ is congruent to $T_1 \Longrightarrow T_2\ R\, T_1$. Hence, R is symmetric.

3. $(T_1 R\, T_2) \wedge (T_2 R\, T_3) \Longrightarrow T_1$ is congruent to T_2 and T_2 is congruent to $T_3 \Longrightarrow T_1$ is congruent to T_3. Hence, R is transitive.

Therefore, R is an equivalence relation.

Exercise: Consider the binary relation R defined over the set $\mathbb{Z}$:

$$\forall\, x,\ y \ \in \mathbb{Z}: \quad x\ R\, y \quad \Longleftrightarrow \quad x + y \text{ is even.}$$

Show that R is an equivalence relation.

Solution:

(a) R is **reflexive** $\Leftrightarrow \forall x \in \mathbb{Z}:\ x\ R\, x$.

1. Let $x \in \mathbb{Z}$, the sum $x + x$ can be written as $2k$ for some integer k (namely, x), so $x + x$ is even. Then $x\ R\ x$ holds, as required.

(b) R is **symmetric** $\Leftrightarrow$ $\forall x, y \in \mathbb{Z} : x\ R\ y \Rightarrow y\ R\ x.$

Let $x, y \in \mathbb{Z}$,

$$\begin{aligned} x\ R\ y \quad &\Rightarrow \quad \exists k \in \mathbb{Z},\ x + y = 2k \\ &\Rightarrow \quad y + x = x + y = 2k \quad \text{(by Commutative Property of Addition)} \\ &\Rightarrow \quad y + x \text{ is even} \\ &\Rightarrow \quad y\ R\ x, \text{ as required} \end{aligned}$$

(c) R is **transitive** $\Leftrightarrow$ $\forall x, y, z \in \mathbb{Z} : (x\ R\ y) \wedge (y\ R\ z) \Longrightarrow x\ R\ z.$

Let $x, y, z \in \mathbb{Z}$,

$$\begin{aligned} (x\ R\ y) \wedge (y\ R\ z) \quad &\Rightarrow \quad \exists\ k, k^{'} \in \mathbb{Z},\ (x + y = 2k) \wedge \left(y + z = 2k^{'}\right) \\ &\Rightarrow \quad x + y + y + z = 2k + 2k^{'} \\ &\Rightarrow \quad x + z = 2k + 2k^{'} - 2y = 2k^{''} / \ k^{''} = k + k^{'} + y \\ &\Rightarrow \quad x + z \text{ is even} \\ &\Rightarrow \quad x\ R\ z, \text{ holds, as required.} \end{aligned}$$

Therefore, R is an equivalence relation.

2.2.3 Equivalences and Partitions

Definition (Equivalence Relation)

Given a partition A_1, A_2, A_3, $\ldots$ of a set A, two elements x and y of A are said to be **equivalent**, with respect to that partition, if they belong to the same subset specified by the partition.

Example: The days of the year are partitioned by seven disjoint sets given by the weekday names of the days. For instance, August 1, 1966, and June 30, 2003, are equivalent in this context since they both belong to the subset called "Monday."

Example: Assuming two words of the English language to be equivalent if they each possess the same number of vowels is an equivalence relation on the set of all words.

Definition (Equivalence Classes)

Given an equivalence relation R over a set A, for any $x \in A$, the equivalence class of x is the set

$$[x] = \{y,\ x\ R\ y\}$$

$[x]$ is the set of all elements of A that are related to x by relation R.

Property: If R is an equivalence relation over A, then every $x \in A$ belongs to exactly one equivalence class.

Theorem 3. If R is an equivalence relation on a set A, then the collection of its equivalence classes is a partition of A. Conversely, if P is a partition of A, then the relation defined by

$$x\ R\ y \Leftrightarrow \exists\ S \in P : x, y \in S$$

is an equivalence relation, and its equivalence classes are the elements of the partition.

Exercise: Provide a proof.

2.2.4 Order Relation

The notion of order relation on a set aims to define the intuitive idea that an object ”precedes” another, ”come before” it, according to a certain criterion of ordering, of disposition of the objects in question.

Definition (Order Relation)

A binary relation R on a set A is called an **order relation** if it is *reflexive, anti-symmetric, and transitive.*

Examples:

1. The “*less-than-or-equal-to”* relation on the set of integers $\mathbb{Z}$ is an order relation.

2. The “*strictly-less-than”* and “*proper-subset*” relations are not order relation because they are not reflexive.

Definition (Total Ordering Relation)

An **order relation** R on A is called a **total ordering** if it satisfies one additional proposition:

$$\boxed{\forall x \in A, \forall y \in A: \ (x\ R\ y) \vee (y\ R\ x)}$$

Examples:

1. The relation $(\mathbb{R}, \leqslant)$ is a total order relation.

2. The relation $(P(E), \subseteq)$ is not a total order relation because $\exists\ A, B \in P(E) : (A \nsubseteq B) \wedge (B \nsubseteq A)$.

Example: Show that the relation "Divides" defined on $\mathbb{N}^*$ is an order relation.

Solution:

1) R is **reflexive** $\Leftrightarrow \forall x \in \mathbb{N}^* :\ x\ R\ x$

We have x divides x, $\forall x \in \mathbb{N}^*$. Therefore, relation "*Divides*" is reflexive.

2) R is **anti-symmetric** $\Leftrightarrow\ \forall x, y \in \mathbb{N}^* : (x\ R\ y) \wedge (y\ R\ x) \Longrightarrow x = y$

Let $x, y \in \mathbb{N}^*$,

$$\begin{aligned}
(x\ R\ y) \wedge (y\ R\ x) \quad &\Rightarrow \quad (x \text{ divides } y) \text{ and } (y \text{ divides } x) \\
&\Rightarrow \quad \exists\ k, k' \in \mathbb{N}^*, \left(y = k' x\right) \wedge (\ x = ky) \\
&\Rightarrow \quad x = kk' x \\
&\Rightarrow \quad kk' = 1 \\
&\Rightarrow \quad k = k' = 1 \in \mathbb{N}^* \\
&\Rightarrow \quad x = y
\end{aligned}$$

So, the relation is anti-symmetric.

3) R is **transitive** $\Leftrightarrow\ \forall x, y, z \in \mathbb{N}^* : (x\ R\ y) \wedge (y\ R\ z) \Longrightarrow x\ R\ z$.

Let $x, y, z \in \mathbb{N}^*$,

$$
\begin{aligned}
(x\ R\ y) \wedge (y\ R\ z) \quad &\Rightarrow \quad \exists\, k, k^{'} \in \mathbb{N}^{*}, (y = k\ x) \wedge \Big(z = k^{'} y \Big) \\
&\Rightarrow \qquad\qquad z = k^{'} k\ x \\
&\Rightarrow \qquad z = k^{"}\ x\ /\ k^{"} = k^{'} k \in \mathbb{N}^{*} \\
&\Rightarrow \qquad x \text{ divides } z \Rightarrow x\ R\ z
\end{aligned}
$$

Hence, the relation is transitive.

Thus, the relation R being reflexive, anti-symmetric and transitive, the relation "*divides*" is an order relation.

2.3 Applications

2.3.1 Functional Relation

The concept of a *function* is one of the most basic mathematical ideas and enters into almost every mathematical discussion.We focus on the concept of *Functional relation* which is called an *application*. We will not study derivatives or integrals, but rather the notions of *injective and surjective applications*, how to compose *applications*, and when they are *invertible*.

Definition (Function)

Let E and F be sets. A **function** is a relation from a set E to another set F, denoted by $f : E \to F$, that *every element* $x \in E$ assigns *at most a unique element* $y \in F$ satisfying $x\ f\ y$. To indicate this relation between x and y we usually write $y = f(x)$.

$$\boxed{x\ f\ y \Longleftrightarrow y = f(x)}$$

We say that:

- y is the **image** of x (under f).
- x is the **pre-image** of y (under f).
- f **maps** x onto y, and symbolize this statement by $x \overset{f}{\longmapsto} y$.
- E is the **starting set** of f.

- F is the **arrival set** or **codomain** of f.

- G_f is the **graph** of the function f, given by: $G_f = \{(x, y) \in E \times F \ / \ y = f(x)\}$

Note that

$$\begin{array}{rccc} f: & E & \longrightarrow & F \\ & x & \longmapsto & y \end{array}$$

Definition (Domain Definition)

The **Domain** D_f of a function f of E in F is the set of elements $x \in E$ satisfying: there is one and only one element $y \in F$ such that $y = f(x)$.

Definition (Range)

We call **range** of a function f the subset of F with *preimages*.

Definition (Application)

An **application** f is a function of E in F whose domain definition D_f is equal to E.

Definition (Application)

An **application** from a set E to a another set F is a relation which to *every element* $x \in E$ assigns a *unique element* $y \in F$. Formally, using predicate logic:

$$\boxed{f \ \textbf{Application} \Leftrightarrow \begin{cases} 1) & \forall\, x \in E,\ \exists\, y \in F: \quad y = f(x) \\ 2) & \forall x_1,\ x_2 \in E : x_1 = x_2 \Rightarrow f(x_1) = f(x_2) \end{cases}}$$

we can write also,

$$\boxed{f \ \textbf{Application} \Leftrightarrow \forall\, x \in E,\ \exists!\, y \in F: \quad y = f(x)}$$

An **application** (or function) is a triplet $f = (E, F, R)$, where E and F are two sets and $G_f \subseteq E \times F$ is a functional relation (See Figure 2.9).



Figure 2.9

Example: Consider the assignment rule f **:** $E = \{1, 2, 3, 4\} \to F = \{x, y, z\}$ which is defined by:

1. $G = \{(1, x), (2, y), (3, z), (4, x)\}$

2. $G = \{(1, x), (2, x), (3, x), (4, y)\}$

3. $G = \{(1, z), (2, y), (3, x)\}$

4. $G = \{(1, y), (2, x), (3, y), (3, z), (4, x)\}$

The first two relations are applications and the third relation is function with D_f is $\{1, 2, 3\}$ but not the last one.

2.3.2 Equality - Extension - Restriction

Definition (The equality of applications)

Two applications f and g are called equal if and only if they have the same domain E, the same codomain F and the same graphic $G_f = Gg$. If $f, g : E \longrightarrow F$, the **equality** $f = g$ is equivalent to $f(x) = g(x), \forall x \in E$, that is to say:

$$\boxed{f = g \Longleftrightarrow \forall x \in E,\ f(x) = g(x)}$$

Definition (Extension of an application-Restriction of an application)

Let $f : E \to F$ be an application and X and Y be sets such that $E \subseteq X$ and $F \subseteq Y$. An **extension** of f to X is an application $g : X \to Y$ such that $f(x) = g(x)$ for all $x \in E$. Alternatively, g is an extension of f to X if f is the **restriction** of g to E.

2.3.3 Image and Inverse Image of a Subsets

Often in mathematics, particularly in analysis and topology, one is interested in finding the set of image points or inverse image of an application acting on a given set, which brings us to the two following definitions that are waiting to be understood.

Definition (Image of a Subset)

Let $f : E \longrightarrow F$ and consider the subset $A \subset E$. The **image** of the subset A under f , which we write $f(A)$, is the subset of F that consists of the *images* of the elements of A (See Figure 2.10)

$$\boxed{f(A) = \{f(x),\ \ x \in A\}}$$

$$\boxed{y \in f(A) \Longleftrightarrow \exists\, x \in A, y = f(x)}$$

Definition (Inverse Image of a Subset)

Let $f : E \longrightarrow F$ and consider the subset $B \subset F$. The **inverse image** of the subset B under f, which we write $f^{-1}(B)$ is the subset of E that consists of the *pre-images* of elements in B (See Figure 2.10)

$$\boxed{f^{-1}(B) = \{x,\ f(x) \in B\}}$$

Example: Let $E = \{1, 2, 3, 4\}$ and $F = \{a, b, c\}$ and define an application $f : E \longrightarrow F$ such that $f(1) = f(2) = a$, $f(3) = f(4) = c$. Let $A \subset E$, $A = \{1, 2, 3\}$. Then $f(A) = \{a, c\}$. Also for example $f^{-1}(\{b\}) = \emptyset$, $f^{-1}(\{a, c\}) = A$, $f^{-1}(\{b, c\}) = \{3\}$.



Figure 2.10

Properties

Given an application $f : E \longrightarrow F$ where A, B, are subsets of E and C, D, are subsets of F, we have the following properties. Notice how the inverse image always preserves unions and intersections, although not always true for the image of an application. Then the images of intersections and unions satisfy:

1. $f(A \cap B) \subseteq f(A) \cap f(B)$.

2. $f(A \cup B) = f(A) \cup f(B)$.

3. $f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$.

4. $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$.

5. $A \subseteq B \Longrightarrow f(A) \subseteq f(B)$.

6. $C \subseteq D \Longrightarrow f^{-1}(C) \subseteq f^{-1}(D)$.

7. $f^{-1}(\overline{C}) = \overline{f^{-1}(C)}$.

Exercise: Let $f(x) = 1 + x^2$. Find the following:

1. $f(\{-1, 1\})$.

2. $f([-2, 2])$.

3. $f([-2, 3])$.

4. $f^{-1}(\{1, 5, 10\})$.

5. $f^{-1}([0,1])$.

6. $f^{-1}([2,5])$.

Solution:

1. $f(\{-1,1\}) = \{2\}$.

2. $f([-2,2]) = f([-2,0] \cup [0,2]) = f([-2,0]) \cup f([0,2]) = [1,5]$.

3. $f([-2,3]) = f([-2,0] \cup [0,3]) = f([-2,0]) \cup f([0,3]) = [1,10]$.

4. $f^{-1}(\{1,5,10\}) = \{0,2,-2,3,-3\}$.

5. $f^{-1}([0,1]) = \{0\}$.

6. $f^{-1}([2,5]) = [-2,-1] \cup [1,2]$.

2.3.4 Injective, Surjective and Bijective Applications

Definition (Injective)

An **injective** application (or **one-to-one** application) $f : E \longrightarrow F$, is an application for which *every element of the range of the application corresponds to exactly one element of the domain.* Formally:

$$\boxed{f \ \textbf{Injective} \Longleftrightarrow \begin{cases} \forall x_1,\ x_2 \in E : x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2) \\ \text{or} \\ \forall x_1,\ x_2 \in E : f(x_1) = f(x_2) \Rightarrow x_1 = x_2 \end{cases}}$$

In words, this says that all elements in the domain of f have different images (See Figure 2.11)

Example. Consider an application $f : \mathbb{R} \longrightarrow \mathbb{R}$, $f(x) = 4x - 1$. We want to know whether each element of $\mathbb{R}$ has a different image. In fact, this function is a line, so one may "see" that two distinct elements have distinct images, but let us try a proof of this.



Figure 2.11

$$
\begin{aligned}
f(x_1) = f(x_2) \quad &\Longrightarrow \quad 4x_1 - 1 = 4x_2 - 1 \\
&\Longrightarrow \qquad 4x_1 = 4x_2 \\
&\Longrightarrow \qquad x_1 = x_2
\end{aligned}
$$

Therefore f is injective.

Example. Consider an application $g : \mathbb{R} \longrightarrow \mathbb{R}$, $g(x) = x^2$. A property of injectivity of g is not true by providing an example where it does not hold. The two elements $x_1 = 1$ and $x_2 = -1$ are both sent to $g(x_1) = g(x_2) = 1$.

The other definition that always comes in pair with that of injective is that of surjective.

Definition (Surjective)

An application $f : E \longrightarrow F$ is **surjective** (or **onto**) if and only if *for every element* $y \in F$, *there is an element* $x \in E$ with $y = f(x)$:

$$\boxed{\forall y \in F, \exists\, x \in E : y = f(x)}$$

In words, each element in the co-domain of f has a pre-image (See Figure 2.12)

Example. Consider again $f : \mathbb{R} \longrightarrow \mathbb{R}$, $f(x) = 4x - 1$. We want to know whether each element of $\mathbb{R}$ has a preimage.



Figure 2.12

$$f(x) = y \implies 4x - 1 = y$$

$$\implies 4x = y + 1$$

$$\implies x = \frac{y+1}{4} \in \mathbb{R}$$

Therefore f is surjective.

Example. Consider again $g : \mathbb{R} \longrightarrow \mathbb{R}$, $g(x) = x^2$. A property of surjectivity of g is not even true by providing an example where it does not hold. If $y = -1$, there is no $x \in \mathbb{R}$ such that $g(x) = x^2 = -1$.

Exercise: The function f is defined by: $f : \mathbb{R} \to \mathbb{R} : x \mapsto x^2 - 6x$

1. Give an example to show that f is not injective.

2. Give an example to show that f is not surjective.

Solution:

1. $f(6) = f(0) = 0$ but $6 \neq 0$, therefore the application is not injective.

2. $f(x) = x^2 - 6x = (x - 3)^2 - 9$

let $y = -10$ then $f(x) = -10 \implies (x-3)^2 - 9 = -10$

$$\implies \quad (x-3)^2 = -1$$

There is no real number, x such that $f(x) = -10$ the application is not surjective.

Or the range of the application is $y \geq 2$. The range of the function is not $\mathbb{R}$ (the codomain), therefore the application is not surjective.

We next combine the definitions of an application which is injective and surjective, to get:

Definition 1 (Bijective)

An application $f : E \longrightarrow F$ is **bijective** if and only if *it is both injective and surjective* (See Figure 2.13)

Definition 2 (Bijective)

An application $f : E \longrightarrow F$ is **bijective** if and only if *for every element $y \in F$, there is a unique element $x \in E$* with $y = f(x)$:

$$\boxed{\forall y \in F, \exists!\, x \in E : y = f(x)}$$



Figure 2.13

Example: Consider the application $f : \mathbb{R} \longrightarrow \mathbb{R}$, $f(x) = 4x - 1$, which we have just studied in two previous examples. We know it is both injective and surjective, therefore it is a bijection.

Bijections have a special feature: they are **invertible**, formally:

Definition 1 (Inverse Application)

Let $f : E \longrightarrow F$ be a bijection. Then the **inverse application** of f , $f^{-1} : F \longrightarrow E$ is defined elementwise by: $f^{-1}(y)$ is the *unique element* $x \in E$ such that $f(x) = y$. We say that f is **invertible**.

Example: Let us consider again at our two previous examples, namely, $f(x) = 4x - 1$ and $g(x) = x^2$. Then, the application g is not a bijection, so it cannot have an inverse. Now f is an application bijective, so we can compute its inverse.

$$\begin{aligned} y = f(x) \iff & \quad y = 4x - 1 \\ \iff & \quad y + 1 = 4x \\ \iff & \quad x = \tfrac{y+1}{4} \\ \iff & \quad f^{-1}(y) = \tfrac{y+1}{4} \end{aligned}$$

Property: Let $f : E \longrightarrow F$ be a bijective application , then $f^{-1} : F \longrightarrow E$ is a bijective application.

2.3.5 Examples of Applications

- **Identity Application**. Let A be a set; by the identity application on A we mean the application $I_A : A \to A$ given by

$$\boxed{I_A(x) = x}$$

  * I_A is injective, $I_A(x) = I_A(y) \Longrightarrow x = y$ $(I_A(x) = x$ and $I_A(y) = y)$; thus the injection holds.

  * I_A is surjective because, obviously, the *range* of I_A is A.

  * Thus I_A is bijective.

- **Constant Application**. Let A and B be sets, and let b be an element of B. By the constant application f_b we mean the application $f_b : A \to B$ given by:

$$\boxed{f_b(x) = b, \forall x \in A}$$

- **Characteristic Application.** The characteristic application of a set is used to solve some difficult problems of set theory found in undergraduate studies. Let's consider $A \subset E \neq \emptyset$ (a universal set), then $f_A : E \to \{0, 1\}$, where the application

$$f_A(x) = \begin{cases} 1, & x \in A \\ \\ 0, & x \notin A \end{cases}$$

is called the characteristic application of the set A.

2.3.6 Operations on Applications

Given two applications, it may be possible to combine them in different ways to create more complicated applications. If the domains and codomains of the two applications agree and if the codomain supports arithmetic, we may define arithmetic operations on the applications by point-wise operations on their images.

Arithmetic Operations on Applications

Let $f : E \to F$ and $g : E \to F$ be two applications sharing a common domain E and let α be a real number. Then $f + g$, $f - g$, $\alpha \cdot f$, $f \cdot g$, and f/g $(g(x) \neq 0)$ denote the following applications from E to F :

$(f + g)(x) = f(x) + g(x)$.

$(f - g)(x) = f(x) - g(x)$.

$(\alpha \cdot f)\ (x) = \alpha \cdot f(x)$.

$(f \cdot g)(x) = f(x) \cdot g(x)$.

$(f/g)(x) = f(x)/g(x)$, provided $g(x) \neq 0$.

Exercise: Find counter-examples to each of these statements for $f : \mathbb{R} \to \mathbb{R}$ and $g : \mathbb{R} \to \mathbb{R}$:

(a) If f and g are surjective, then $(f + g)$ is surjective.

Suppose $f(x) = x$ and $g(x) = -x$. Then $(f + g)\,(x) = x - x = 0$.

(b) If f and g are surjective, then $f \cdot g$ is surjective.

The same $f(x) = x$ and $g(x) = -x$ from above work; $(f \cdot g)(x) = -x^2$, which is not surjective.

Composition of Applications

In addition to arithmetic operations on applications, there is another operation called composition of applications which is more set-theoretic or algebraic in nature. A composite of two applications is satisfied if the codomain of the first application agrees with the domain of the second.

If $f : E \to F$ and $g : F \to G$ are applications in which the codomain of f equals the domain of g, then the assignment $h(x) = g(f(x))$ defines an application $h : E \to G$. For, given any $x \in E$, there is a unique $y \in F$ such that $y = f(x)$, since f is an application. Similarly, since g is an application, $g(f(x))$ is a unique image in G. Thus each element x from E yields a unique image $z = g(f(x))$ in G, guaranteeing that h is an application from E into G. This legitimizes the following definition.

Definition (Composite Applications)

If $f : E \to F$ and $g : F \to G$, then the composite application f followed by g is the application $g \circ f$ such that:

$$\begin{array}{rccl} g \circ f : & E & \to & G \\ & x & \mapsto & (g \circ f)(x) = g(f(x)) \end{array}$$

Example: If $f(x) = -4x + 9$ and $g(x) = 2x - 7$, find $(f \circ g)(x)$ and $(g \circ f)(x)$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) \\ &= -4g(x) + 9 \\ &= -4(2x - 7) + 9 \\ &= -8x + 28 + 9 \\ &= -8x + 37 \end{aligned}$$

Thus, $(f \circ g)(x) = -8x + 37$.

$$
\begin{aligned}
(g \circ f)(x) &= g(f(x)) \\
&= 2f(x) - 7 \\
&= 2(-4x + 9) - 7 \\
&= -8x + 18 - 7 \\
&= -8x + 11
\end{aligned}
$$

Thus, $(g \circ f)(x) = -8x + 37$.

We remark that $(f \circ g)(x)$ and $(g \circ f)(x)$ produced different answers.

Properties: Suppose f, g, and h are application that can be composed in the order given.

1. Composition is not Commutative: $g \circ f \neq f \circ g$.
2. If f and g are both injective applications, then so is $g \circ f$.
3. If f and g are both surjective applications, then so is $g \circ f$.
4. If f and g are both bijective applications, then so is $g \circ f$.
5. Composition is associative: $(h \circ g) \circ f = h \circ (g \circ f)$.

Definition 2 (Inverse Applications)

If $f : E \to F$ and $g : F \to E$, then f and g are inverse applications of one another relative to composition iff

$$g \circ f = I_E \text{ and } f \circ g = I_F.$$

Example: Show that the application $g : \mathbb{R} \to \mathbb{R}$ defined by $g(x) = \frac{x-1}{2}$ is an inverse for the application $g : \mathbb{R} \to \mathbb{R}$ defined by $f(x) = 2x + 1$.

$$g(f(x)) = \frac{f(x)-1}{2} = \frac{2x+1-1}{2} = x$$

$$f(g(x)) = 2g(x) + 1 = 2\left(\frac{x-1}{2}\right) + 1 = x$$

Thus g is f's inverse.

Properties: Let $f : E \to F$ and $g : F \to G$

1. If f has an **inverse**, then it is unique.

2. The composition $g \circ f$ of two invertible applications f and g is invertible. Moreove , the composition of the inverses in the reverse order

$$\boxed{(g \circ f)^{-1} = f^{-1} \circ g^{-1}}$$

Example: Let an application $f : E \to F$.

Determine the inverse application for $f(x) = \frac{x}{x+1}$. Assume that f is defined for as inclusive a set of real numbers as possible and that the codomain of f is its range.

The equation $y = \frac{x}{x+1}$ defines an application on $E = \mathbb{R} - \{-1\}$.

We will check its codomain after we determine which values y can be. Solving $y = \frac{x}{x+1}$ for x, we get the following:

$$\begin{aligned} y = \tfrac{x}{x+1} \quad &\Rightarrow \quad y\,(x+1) = x \\ &\Rightarrow \quad yx + y = x \\ &\Rightarrow \quad yx - x = -y \\ &\Rightarrow \quad x = \tfrac{y}{1-y} = g(y) \end{aligned}$$

Since there are x-values for all y except $y = 1$, our domain for g and our codomain for f must be taken to be $F = \mathbb{R} - \{1\}$.

For these x- and y-values the above solution process is reversible. The inverse application is therefore given by $f^{-1}(y) = \frac{y}{1-y}$.

2.4 The Inverse Trigonometric Application

In this section, we concern ourselves with finding inverses of the (circular) trigonometric applications. Our immediate problem is that, owing to their periodic nature, none of the circular applications is *injective*. To remedy this, we restrict the domains of the circular applications to obtain an *injective* application.

2.4.1 Arccosine Application

We first consider $f(x) = \cos(x)$. Choosing the interval $[0, \pi]$ allows us to keep the range as $[-1, 1]$ as well as the property of being *bijective*.



Figure 2.14

Recall from Subsection 2.3.4 that the inverse of an application f is typically denoted f^{-1}. For this reason,we can use the notation $f^{-1}(x) = \cos^{-1}(x)$ for the inverse of $f(x) = \cos(x)$ (See Figures 2.14–2.15)

Remark: It is far too easy to confuse $\cos^{-1}(x)$ with $\frac{1}{\cos(x)}$ so we will not use this notation in our text.

Notation: We use the notation $f^{-1}(x) = \arccos(x)$, read "**arc-cosine** of x".



Figure 2.15

Formally:

$$\begin{aligned} f^{-1}: \quad [-1,1] \quad &\to \quad [0,\pi] \\ x \quad &\mapsto \quad f^{-1}(x) \quad = \arccos(x) \end{aligned}$$

We list some important facts about the arccosine applications in the following properties.

Properties

$\arccos(x) = y$ if and only if $y \in [0,\pi]$ and $\cos(y) = x$.

$\cos(\arccos(x)) = x$ provided $x \in [-1,1]$.

$\arccos(\cos(x)) = x$ provided $x \in [0,\pi]$.

2.4.2 Arcsine Application

We restrict $f(x) = \sin(x)$ in a similar manner, although the interval of choice is $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ (See Figure 2.16)

It should be no surprise that we call $f^{-1}(x) = \arcsin(x)$, which is read "**arc-sine** of x"(See Figure 2.17)



Figure 2.16

Formally:

$$\begin{aligned} f^{-1}: \quad [-1,1] \quad &\to \quad \left[-\tfrac{\pi}{2}, \tfrac{\pi}{2}\right] \\ x \quad &\mapsto \quad f^{-1}(x) \quad = \arcsin(x) \end{aligned}$$

We list some important facts about the arcsine applications in the following properties.

Properties

$\arcsin(x) = y$ if and only if $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ and $\sin(y) = x$.

$\sin(\arcsin(x)) = x$ provided $x \in [-1, 1]$.

$\arcsin(\sin(x)) = x$ provided $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.



Figure 2.17

Exercise: Find the exact values of the following.

1) $\arccos(\frac{1}{2})$	5) $\arcsin(\frac{\sqrt{2}}{2})$
2) $\arccos(-\frac{\sqrt{2}}{2})$	6) $\arcsin(-\frac{1}{2})$
3) $\arccos(\cos\left(\frac{\pi}{6}\right))$	7) $\arccos(\cos(\frac{11\pi}{6}))$
4) $\cos\left(\arccos(-\frac{3}{5})\right)$	8) $\sin\left(\arccos(-\frac{3}{5})\right)$

Solution:

1. To find $\arccos(\frac{1}{2})$, we need to find the real number y (or, equivalently, an angle measuring y radians) which verifies $y \in [0, \pi]$ and with $\cos(y) = \frac{1}{2}$. We know $y = \frac{\pi}{3}$ meets these criteria, so $\arccos(\frac{1}{2}) = \frac{\pi}{3}$.

2. The number $y = \arccos(-\frac{\sqrt{2}}{2}) \in [0, \pi]$ with $\cos(y) = -\frac{\sqrt{2}}{2}$. Our answer is $y = \frac{3\pi}{4}$.

3. Since $\frac{\pi}{6} \in [0, \pi]$, we could simply refer to the properties of arccosine applications to get $\arccos(\cos\left(\frac{\pi}{6}\right)) = \frac{\pi}{6}$.

4. One way to simplify $\cos\left(\arccos(-\frac{3}{5})\right)$ is to use the properties of arccosine applications directly. Since $-\frac{3}{5} \in [-1, 1]$, we have $\cos\left(\arccos(-\frac{3}{5})\right) = -\frac{3}{5}$.

5. The value of $\arcsin(\frac{\sqrt{2}}{2})$ is a real number $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ with $\sin(y) = \frac{\sqrt{2}}{2}$. The number we seek is $y = \frac{\pi}{4}$. Hence, $\arcsin(\frac{\sqrt{2}}{2}) = \frac{\pi}{4}$.

6. To find $\arcsin(-\frac{1}{2})$, we seek the number $y \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ with $\sin(y) = -\frac{1}{2}$. The answer is $y = -\frac{\pi}{6}$ so that $\arcsin(-\frac{1}{2}) = -\frac{\pi}{6}$.

7. Since $\frac{11\pi}{6}$ does not fall between 0 and π, the properties of the arcsine applications does not apply. We are forced to work through from the inside out starting with $\arccos(\cos(\frac{11\pi}{6})) = \arccos(\frac{\sqrt{3}}{2})$. We know $\arccos(\frac{\sqrt{3}}{2}) = \frac{\pi}{6}$. Hence, $\arccos(\cos(\frac{11\pi}{6})) = \frac{\pi}{6}$.

8. As in the previous question, we let $y = \arccos(-\frac{3}{5})$ so that $\cos y = -\frac{3}{5}$ for $y \in [0, \pi]$. Since $\cos y < 0$, we can narrow this down a bit and conclude that $\frac{\pi}{2} < y < \pi$. In terms of y, then, we need to find $\sin\left(\arccos(-\frac{3}{5})\right) = \sin y$. Using the Pythagorean Identity $\cos^2 y + \sin^2 y = 1$, we get $\left(-\frac{3}{5}\right)^2 + \sin^2 y = 1$ or $\sin y = \pm\frac{4}{5}$. We choose $\sin y = \frac{4}{5}$. Hence, $\sin\left(\arccos(-\frac{3}{5})\right) = \frac{4}{5}$.

The next pair of application we wish to discuss are the inverses of tangent and cotangent, which are named arctangent and arccotangent, respectively.

2.4.3 Arctangent Application

We restrict $f(x) = \tan(x)$ to its fundamental cycle on $\left]-\frac{\pi}{2}, \frac{\pi}{2}\right[$ to obtain $f^{-1}(x) = \arctan(x)$. Among other things, note that the *vertical asymptotes* $x = -\frac{\pi}{2}$ and $x = \frac{\pi}{2}$ of the graph of $f(x) = \tan(x)$ become the horizontal asymptotes $y = -\frac{\pi}{2}$ and $y = \frac{\pi}{2}$ of the graph of $f^{-1}(x) = \arctan(x)$. We show these graphs on Figure 2.18.

We list some of the basic properties of the arctangent application.

Properties

$\arctan(x) = y$ if and only if $y \in \left]-\frac{\pi}{2}, \frac{\pi}{2}\right[$ and $\tan(y) = x$.

$\tan(\arctan(x)) = x$ provided $x \in \mathbb{R}$.

$\arctan(\tan(x)) = x$ provided $x \in \left]-\frac{\pi}{2}, \frac{\pi}{2}\right[$.



Figure 2.18

2.4.4 Arccotangent Application

We restrict $f(x) = \cot(x)$ to its fundamental cycle on $]0, \pi[$ to obtain $f^{-1}(x) = arccot(x)$. Once again, the *vertical asymptotes* $x = 0$ and $x = \pi$ of the graph of $f(x) = \cot(x)$ become the *horizontal asymptotes* $y = 0$ and $y = \pi$ of the graph of $f^{-1}(x) = arccot(x)$. We show these graphs on Figure 2.19.

We list some of the basic properties of the arccotangent application.

Properties

$arccot(x) = y$ if and only if $y \in]0, \pi[$ and $\cot(y) = x$.

$\cot(arccot(x)) = x$ provided $x \in \mathbb{R}$.

$arccot(\cot(x)) = x$ provided $x \in]0, \pi[$.

Exercise: Find the exact values of the following.

1. $\arctan(\sqrt{3})$.

2. $arccot(-\sqrt{3})$.

3. $\cot(arccot(-5))$



Figure 2.19

Solution:

1. We know $\arctan(\sqrt{3})$ is the real number $y \in \left]-\frac{\pi}{2}, \frac{\pi}{2}\right[$ with $\tan(y) = \sqrt{3}$. We find $y = \frac{\pi}{3}$, so $\arctan(\sqrt{3}) = \frac{\pi}{3}$.

2. The real number $y = arccot(-\sqrt{3}) \in]0, \pi[$ with $\cot(y) = -\sqrt{3}$. We get $arccot(-\sqrt{3}) = \frac{5\pi}{7}$.

3. We can apply properties of the arccotangent application directly and obtain $\cot(arccot(-5)) = -5$.

FAMOUS ISLAMIC MATHEMATICIANS

Ibrahim Ibn Sinan found out multiple **Trigonometry identities**, and was key in finding out about **tangents** in circles. His most famous piece of work is said to be on the quadrature of the parabola

Ibrahim Ibn Sinan

Chapter 3

Real functions of one real variable

Descartes was introducing algebra into geometry. His study to the solutions of equations of degree 3, 4, 5 and more led to the concept of the function.

In this chapter we shall study limit, continuity and differentiability of real valued functions defined on certain sets.

3.1 Elementary Functions

In mathematics, an "*elementary function*" is a function of a single variable composed of particular simple functions.

Basic examples:

The elementary functions of (x) of mathematics comprise:

- *Trigonometric functions:* $x \mapsto \sin x, \cos x, \tan x, \cot x$
- *Exponential functions:* $x \mapsto \exp x$
- *Logarithms:* $x \mapsto \ln x$
- *Inverse trigonometric functions*: $x \mapsto \arcsin x,\ \arccos x, \arctan x, arccotx$

- *Hyperbolic functions* :

$$x \mapsto \cosh x = \frac{\exp(x) + \exp(-x)}{2}, \quad x \mapsto \sinh x = \frac{\exp(x) - \exp(-x)}{2}$$

- *Inverse hyperbolic functions:* $x \mapsto \arg chx, \ x \mapsto \arg shx$.

3.2 Limit of a Function

The notion of a limit is a fundamental concept of calculus. More particularly, limits allow us to look at what happens in a **very**, **very** small region around a point.

Example 1: Values of $f(x) = \frac{x^2-4}{x-2}$ may be computed near $x = 2$

x	1.9	1.99	1.999 $\rightarrow$	$\leftarrow$ 2.001	2.01	2.1
$f(x)$	3.9	3.99	3.999 $\rightarrow$	$\leftarrow$ 4.01	4.01	4.1

$$\lim_{x \to 2} f(x) = 4$$

Definition (Neighbourhood)

For $x_0 \in \mathbb{R}$, an open interval of the form $]x_0 - \delta, x_0 + \delta[$ for some $\delta > 0$ is called a **neighbourhood** of x_0.

3.2.1 Limit of a function at a point

Definition

A real valued function $f : D \to \mathbb{R}$ has "*limit value L as x tends to a finite value x_0*" if one can demonstrate that for any positive number ε (no matter how small), all the values $f(x)$ of the function will eventually be this close to the value L by restricting x to values very close, but not equal, to x_0. That is, one can produce a positive number δ so that if x, different from x_0, lies between $x_0 - \delta$ and $x_0 + \delta$ so then we can be sure that the value

$f(x)$ lies between $L-\varepsilon$ and $L+\varepsilon$. Formally:

$$\forall \varepsilon > 0, \exists\, \delta > 0, \forall x \in D : x \in \,]x_0 - \delta, x_0 + \delta[\, \Rightarrow\ f(x) \in \,]L-\varepsilon, L+\varepsilon[$$

or we can write

$$\forall \varepsilon > 0, \exists\, \delta > 0, \forall x \in D : |x - x_0| < \delta \Rightarrow |\, f(x) - L| < \varepsilon$$

If a function $f(x)$ has a limit value L as x approaches a finite value x_0, we write:

$$\lim_{x \to x_0} f(x) = L$$

Example 1: Show that $\lim\limits_{x \to 4} (2x-1) = 7$. We have $f(x) = 2x-1$, $x_0 = 4$ and $L = 7$ and the question we must answer is "how close should x be to 4 if want to be sure that $f(x) = 2x-1$ differs less than ε from $L = 7$?"

To figure this out we try to get an idea of how big $|\, f(x) - L|$ is:

$|\, f(x) - L| \quad = \quad |(2x-1) - 7| \quad = 2 \cdot |x - 8|$

So, if $2 \cdot |x - x_0| < \varepsilon$ then we have $|\, f(x) - L| < \varepsilon$, i.e.

$|x - x_0| < \frac{\varepsilon}{2} \quad \Longrightarrow \quad |f(x) - L| < \varepsilon.$

We can therefore choose $\delta = \frac{\varepsilon}{2}$. No matter what $\varepsilon > 0$ we are given our δ will also be positive,

$$\forall \varepsilon > 0, \exists\, \delta > 0\ , \forall x \in D\ :\ |x - 4| < \delta \quad \Longrightarrow \quad |(2x-1) - 7| < \varepsilon$$

That shows that $\lim\limits_{x \to 4} f(x) = 7$.

Definitions (Left limit and right limit)

(i) We say that f has the *left limit* $L \in \mathbb{R}$ as x tends to x_0 iff

$$\forall \varepsilon > 0, \exists\, \delta > 0, \forall x \in D : x \in \,]x_0 - \delta, x_0[\, \Rightarrow\ f(x) \in \,]L-\varepsilon, L+\varepsilon[$$

and in that case we write:

$$\lim_{\substack{< \\ x \to x_0}} f(x) = L \quad \text{or} \quad \lim_{x \to x_0^-} f(x) = L$$

(ii) We say that f has the *right limit* $L \in \mathbb{R}$ as x tends to x_0 iff

$$\forall \varepsilon > 0, \exists\, \delta > 0, \forall x \in D : x \in \,]x_0, x_0 + \delta[\Rightarrow \; f(x) \in \,]L - \varepsilon, L + \varepsilon[$$

and in that case we write:

$$\lim_{\substack{> \\ x \to x_0}} f(x) = L \quad \text{or} \quad \lim_{x \to x_0^+} f(x) = L$$

Theorem 3 (Existence of the limit)

Let f be a real valued function defined on a set $D \subset \mathbb{R}$, then $\lim\limits_{x \to x_0} f(x)$ exists if and only if:

$$\lim_{x \to x_0^-} f(x) \; = \lim_{x \to x_0^+} f(x) = \lim_{x \to x_0} f(x)$$

Example 1:

Let $f : [-1, 1] \to \mathbb{R}$ defined by:

$$f(x) = \begin{cases} 0, & -1 \leq x \leq 0 \\ \\ 1, & 0 < x \leq 1 \end{cases}$$

$\lim\limits_{x \to x_0} f(x)$ does not exist because $\lim\limits_{x \to 0^-} f(x) \; = 0$ and $\lim\limits_{x \to 0^+} f(x) = 1$.

Example 2:

Let $f : \mathbb{R} \to \mathbb{R}$ defined by:

$$f(x) = \sin\left(\frac{1}{x}\right)$$

We see in Figure 3.1, that $\sin\left(\frac{1}{x}\right)$ oscillates between $+1$ and -1 as $x \to 0$. This means that $f(x)$ gets close to any number between $+1$ and -1 as $x \to 0$, but that the function $f(x)$ never stays close to any particular value because it keeps oscillating up and down.

Here again, the limit $\lim_{x \to 0} f(x)$ does not exist.



Figure 3.1

3.2.2 Limit of a function at infinity

Definition

A function f has limit value L as x becomes large if one can demonstrate that for any positive number ε there exists a positive number A, such that all the values $f(x)$ of the function lies between $L - \varepsilon$ and $L + \varepsilon$ for $x > A$. We write: $\lim_{x \to +\infty} f(x) = L$. One can similarly define the notion of a limit as x becomes large and negative: $\lim_{x \to -\infty} f(x) = L$.

Example 1: Let's compute

$$\lim_{x \to \infty} \frac{5x^2 + 1}{4x^2 + 3x - 2}$$

We divide the numerator and denominator by x^2, and you get

$$\lim_{x \to \infty} \frac{5x^2 + 1}{4x^2 + 3x - 2} = \lim_{x \to \infty} \frac{5 + \frac{1}{x^2}}{4 + \frac{3}{x} - \frac{2}{x^2}} = \frac{5}{4}.$$

Example 2: Compute

$$\lim_{x \to \infty} \frac{x}{x^5 - 2}$$

We divide numerator and denominator by x^5. This leads to

$$\lim_{x\to\infty} \frac{\frac{1}{x^4}}{1-\frac{2}{x^5}} = \frac{0}{1} = 0.$$

3.2.3 Properties of the limit

The following properties remain true if one replaces each limit by a one-sided limit, or a limit for $x \to \infty$.

Let f and g be two given functions whose limits for $x \to x_0$ we know,

$$\lim_{x\to x_0} f(x) = L_1, \lim_{x\to x_0} g(x) = L_2.$$

Then:

1. $\lim\limits_{x\to x_0} (f+g)(x) = L_1 + L_2.$
2. $\lim\limits_{x\to x_0} (f\cdot g)(x) = L_1 \cdot L_2.$
3. $\lim\limits_{x\to x_0} (\lambda\cdot f)(x) = \lambda \cdot L_1.$
4. $\lim\limits_{x\to x_0} \frac{f(x)}{g(x)} = \frac{L_1}{L_2}$, if $\lim\limits_{x\to x_0} g(x) \neq 0.$

Theorem 4. Suppose that

$$f(x) \leq g(x) \leq h(x)$$

(for all x) and that

$$\lim_{x\to x_0} f(x) = \lim_{x\to x_0} h(x)$$

Then

$$\lim_{x\to x_0} f(x) = \lim_{x\to x_0} g(x) = \lim_{x\to x_0} h(x)$$

Corollary.

If $\lim\limits_{x\to x_0} f(x) = 0$ and g is a bounded function. Then

$$\lim_{x\to x_0} (f\cdot g)(x) = 0$$

Example 3: Compute

$$\lim_{x\to 0} x^2 \sin\left(\frac{1}{x}\right)$$

Since we have $-1 \leq \sin\left(\frac{1}{x}\right) \leq +1$. Then

$$-x^2 \leq x^2 \cdot \sin\left(\frac{1}{x}\right) \leq +x^2,$$

Since

$$\lim_{x\to 0} -x^2 = \lim_{x\to 0} +x^2 = 0$$

The corollary tells us that

$$\lim_{x\to 0} x^2 \sin\left(\frac{1}{x}\right) = 0$$

3.2.4 Indeterminate Forms

Definition. A function f is said to have an indeterminate form at x_0(where x_0 can be finite or infinite) if:

1. f is continuous on an interval including x_0, except possibly at x_0.

2. When we try to evaluate f at x_0 we obtain one of the following forms:

$$\frac{0}{0}, \frac{\infty}{\infty}, 0 \cdot \infty, \infty - \infty, 1^\infty, \infty^0, 0^\infty$$

Here 0 and 1 represent **variable** quantities approaching the respective value, **NOT** constants with that value. Some indeterminate forms can be solved by rewriting the limit in an equivalent form by factoring through elimination, multiplying by the conjugate, by the trigonometric identities or using L'Hôpital's rule (See the next section).

Example 1: By foctoring

$$\begin{aligned}
\lim_{x\to -1}\tfrac{x^2-x-2}{x^2-2x-3} &= \tfrac{0}{0}\\
\lim_{x\to -1}\tfrac{x^2-x-2}{x^2-2x-3} &= \lim_{x\to -1}\tfrac{(x-2)(x+1)}{(x-3)(x+1)}\\
&= \lim_{x\to -1}\tfrac{(x-2)}{(x-3)}\\
&= \tfrac{3}{4}
\end{aligned}$$

Example 2: By the conjugate

$$\begin{aligned}
\lim_{x\to 4}\tfrac{\sqrt{x}-2}{x-4} &= \tfrac{0}{0}\\
\lim_{x\to 4}\tfrac{\sqrt{x}-2}{x-4} &= \lim_{x\to 4}\tfrac{\left(\sqrt{x}-2\right)\left(\sqrt{x}+2\right)}{(x-4)\left(\sqrt{x}+2\right)}\\
&= \lim_{x\to 4}\tfrac{(x-4)}{(x-4)\left(\sqrt{x}+2\right)}\\
&= \lim_{x\to 4}\tfrac{1}{\left(\sqrt{x}+2\right)}\\
&= \tfrac{1}{4}
\end{aligned}$$

Example 3: By trigonometric identities

$$\begin{aligned}
\lim_{x\to 0}\tfrac{\sin(x)}{\sin(2x)} &= \tfrac{0}{0}\\
\lim_{x\to 0}\tfrac{\sin(x)}{\sin(2x)} &= \lim_{x\to 0}\tfrac{\sin(x)}{2\sin(x)\cos(x)}\\
&= \lim_{x\to 0}\tfrac{1}{2\cos(x)}\\
&= \tfrac{1}{2}
\end{aligned}$$

3.3 Continuous Functions

3.3.1 Continuity of a function at a point

Definitions.

- Let f be a real function on a subset of the real numbers. Then f is **continuous** at x_0 if

$$\begin{cases} 1) & x_0 \in D_f \\ 2) & \lim\limits_{x \to x_0} f(x) = f(x_0) \end{cases}$$

- In particular, if the *left hand limit*, *right hand limit* and the value of the function at $x = x_0$ exist and are equal to each other, i.e.,

$$\lim_{x \to x_0^-} f(x) = \lim_{x \to x_0^+} f(x) = f(x_0)$$

then f is said to be **continuous** at $x = x_0$.

- A function is **continuous** if it is continuous at every x_0 in its domain D_f.

- If it is not continuous there, i.e. if either the limit does not exist or is not equal to $f(x_0)$ we will say that the function is **discontinuous** at x_0.

Example 1: Consider the function

$$f(x) = |x| = \begin{cases} x, & x \geq 0 \\ -x, & x \end{cases}$$

This function is continuous at all x_0, $\lim\limits_{x \to x_0} f(x) = \lim\limits_{x \to x_0} |x| = |x_0| = f(x_0)$.

Example 2:

Let $f : \mathbb{R} \to \mathbb{R}$ defined by:

$$f(x) = \begin{cases} \frac{\sin x}{x}, & x \neq 0 \\ \\ 1, & x = 0 \end{cases}$$

Then f is continuous at $x = 0$, $\lim\limits_{x \to 0} f(x) = \lim\limits_{x \to 0} \frac{\sin x}{x} = f(0) = 1$.

3.3.2 Continuity of a function in an interval

Definition.

1. f is said to be continuous in an open interval $]a, b[$ if it is continuous at every point x_0 in this interval.

2. f is said to be continuous in the closed interval $[a, b]$ if

   - f is continuous in $]a, b[$.
   - f is right continuous at a point a , i.e. $\lim\limits_{x \to a^+} f(x) = f(a)$.
   - f is left continuous at a point b , i.e. $\lim\limits_{x \to b^-} f(x) = f(b)$.

Property. All polynomials, rational functions, trigonometric functions, the inverse trigonometric functions, the absolute value function, the exponential and logarithm functions are continuous everywhere within its domain.

Example : The function $y = \frac{1}{x^2}$ is continuous for $x > 1$ or $x < -1$ but is not continuous on the interval $-1 < x < 1$ (See Figure 3.2).



Figure 3.2

3.3.3 Continuous Extension at a point

We can redefine functions with removable discontinuities to obtain continuous functions.

Proposition.

Let I be an interval , and $x_0 \in I$. Let f be defined on $I - \{x_0\}$ such that $\lim\limits_{x \to x_0} f(x) = l \in \mathbb{R}$.

Consider the function $\widetilde{f}$:

$$\widetilde{f}(x) = \begin{cases} f(x) & \text{for} \quad x \in I - \{x_0\} \\ \\ l & \text{for} \quad x = x_0 \end{cases}$$

then, the function $\widetilde{f}$ is a continuous at x_0.

Example 1: Find a continuous extension of the function $f(x) = \frac{\sin x}{x}$.

The domain of f is $D_f = \mathbb{R}^*$, then f is discontinuous at $x = 0$ because $f(0)$ is not defined. Since $\lim\limits_{x \to 0} f(x)$ exists, the discontinuity is removable.

We know that $\lim\limits_{x \to 0} \frac{\sin x}{x} = 1$. For the function to be continuous at zero we need to define $f(0)$ we make $f(0) = \lim\limits_{x \to 0} \widetilde{f}(x) = \lim\limits_{x \to 0} \frac{\sin x}{x} = 1$.

and redefine the function: $\widetilde{f}(x) = \begin{cases} \frac{\sin x}{x} & \text{for} \quad x \neq 0. \\ \\ 1 & \text{for} \quad x = 0. \end{cases}$

We say $\widetilde{f}$ is the continuous extension of f to $x = 0$.

3.3.4 The Intermediate Value Theorem

It is said that a function is continuous if you can draw its graph without taking your pencil o the paper.

A more precise version of this statement is the intermediate value theorem:

Theorem 5. *If a function f is continuous on a closed interval $[a, b]$, and if y_0 is some number between $f(a)$ and $f(b)$, then there is a number x_0 in the interval $[a, b]$ such that $f(x_0) = y_0$* (See Figure 3.3).



Figure 3.3

Example: Use the Intermediate Value Theorem to prove $x^2 = 2$ has a root.

Consider the function $f : \mathbb{R} \to \mathbb{R}$ defined by: $f(x) = x^2$.

The function f is continuous on a closed interval $[1, 2]$.

One has $f(1) = 2$ and $f(2) = 4$. Since $f(1) \leq 2 \leq f(2)$, the intermediate value theorem with $a = 1$, $b = 2$, $y_0 = 2$ tells us that there is a number x_0 between 1 and 2 such that $f(x_0) = 2$, i.e. for which $x_0^2 = 2$. So the theorem tells us that the *square root* of 2 exists.

3.3.5 Continuity of composite functions

Definition.

Let f and g be real valued functions such that $(f \circ g)$ is defined at x_0. If g is continuous at x_0 and f is continuous at $g(x_0)$, then $(f \circ g)$ is continuous at x_0.

Example: Since both $f(x) = x^2 + 1$ and $g(x) = \cos x$ are continuous on $\mathbb{R}$.

Therefore, both

$(f \circ g)(x) = \cos^2 x + 1$, and

$(g \circ f)(x) = \cos(x^2 + 1)$

are continuous on $\mathbb{R}$.

3.3.6 Continuity of the algebraic combinations of functions

Definition.

If f and g are both continuous at x_0 and λ is any constant, then each of the following functions is also continuous at x_0: The sum $f + g$, the difference $f - g$, the constant multiple λf , the product $f \cdot g$, the quotient f/g , if $g(x_0) \neq 0$, the absolute value $|f|$.

Example: Let $f : \mathbb{R} \to \mathbb{R}$ defined by:

$$f(x) = \begin{cases} x \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Here is a continuous function on $\mathbb{R}$ because

- the inverse function of x , $x \mapsto \frac{1}{x}$ is a continuous on $\mathbb{R}^*$.
- the sine function $x \mapsto \sin x$ is a continuous on $\mathbb{R}$.
- the composite functions $x \mapsto \sin\left(\frac{1}{x}\right)$ is a continuous on $\mathbb{R}^*$
- the function $x \mapsto x$ is a continuous on $\mathbb{R}$.

- the product function $x \mapsto x \sin\left(\frac{1}{x}\right)$ is a continuous on $\mathbb{R}^*$.

- since $\lim\limits_{x \to 0} x \sin\left(\frac{1}{x}\right) = f(0)$, then the function f is a continuous at $x_0 = 0$.

3.4 Differentiability of Functions

Differential calculus is a branch of CALCULUS deals with notions of SLOPE, rates of change and ratios of change. For example, a study of VELOCITY, which can be described as the rate of change of position, falls under the study of differential calculus, as do other concepts that arise in the study of motion.

3.4.1 Differentiability of a function at a point

Definition. **(Differentiability)**

Let f be a real valued function . Then f is said to be **differentiable** at $x_0 \in D_f$ if

$$\lim_{x \to x_0} \frac{f(x) - f(x_0)}{x - x_0} = l \in \mathbb{R}$$

in that case the value l is called the derivative of f at x_0.

The derivative of f at x_0, if exists, is denoted by $f^{'}(x_0)$. Which is read as "f prime of x."

Example 1: The function $f : x \longmapsto \sqrt{x}$ is differentiable at $x_0 = 1$.

$$\lim_{x \to x_0} \frac{f(x)-f(x_0)}{x-x_0} = \lim_{x \to 1} \frac{\sqrt{x}-1}{x-1} = \frac{0}{0}$$

$$\begin{aligned}
\lim_{x \to 1} \frac{\sqrt{x}-1}{x-1} &= \lim_{x \to 1} \frac{\left(\sqrt{x}-1\right)\left(\sqrt{x}+1\right)}{(x-1)\left(\sqrt{x}+1\right)} \\
&= \lim_{x \to 1} \frac{x-1}{(x-1)\left(\sqrt{x}+1\right)} \\
&= \lim_{x \to 1} \frac{1}{\left(\sqrt{x}+1\right)} = \frac{1}{2}
\end{aligned}$$

Since f is differentiable at $x_0 = 1$, then $f^{'}(1) = \frac{1}{2}$.

Example 2: The function $f : x \longmapsto \frac{1}{x}$ is differentiable at $x_0 = 2$.

$$\lim_{x \to x_0} \frac{f(x)-f(x_0)}{x-x_0} = \lim_{x \to 2} \frac{\frac{1}{x}-\frac{1}{2}}{x-2} = \frac{0}{0}$$

$$\lim_{x\to 2}\frac{\frac{1}{x}-\frac{1}{2}}{x-2} = \lim_{x\to 2}\frac{\frac{2-x}{2x}}{x-2}$$

$$= \lim_{x\to 2}\frac{-1}{2x} = -\frac{1}{4}$$

Since f is differentiable at $x_0 = 2$, then $f^{'}(2) = -\frac{1}{4}$.

Definitions (Left Differentiability and Right Differentiability)

- f is left differentiable at a point x_0 , i.e. $\lim\limits_{x\to x_0^-}\frac{f(x)-f(x_0)}{x-x_0} = f_L^{'}(x_0)$.
- f is right differentiable at a point x_0 , i.e. $\lim\limits_{x\to x_0^+}\frac{f(x)-f(x_0)}{x-x_0} = f_R^{'}(x_0)$.

Property. f is differentiable at a point x_0 iff f is left differentiable and right differentiable at this point, i.e. $f_L^{'}(x_0) = f_R^{'}(x_0)$.

Remark. If a quantity y_0 is a FUNCTION of another quantity x_0, $y_0 = f(x_0)$ say, then each change in the x_0-variable, $x_0 \to x_0 + h$, produces a corresponding change in the y_0-variable: $f(x_0) \to f(x_0 + h)$. The ratio of the changes of the two variables is:

$$\frac{f(x_0+h)-f(x_0)}{h}$$

Graphically, this represents the slope of the line segment connecting the two points $(x_0, f(x_0))$ and $(x_0 + h, f(x+h))$ on the graph of the curve $y = f(x)$.

Definition.

The function f is said to be **differentiable** at $x_0 \in D_f$ iff

$$\lim_{h\to 0}\frac{f(x_0+h)-f(x_0)}{h} = f^{'}(x_0)$$

$f^{'}(x_0)$ represents the slope of the (tangent line to the) graph $y = f(x)$ at position x_0, or, alternatively, the instantaneous rate of change of the variable $y = f(x)$ at x_0.

Example 3: The derivative of $f(x) = x^2$ is $f^{'}(x) = 2x$.

$$\lim_{h\to 0}\frac{f(x+h)-f(x)}{h} = \lim_{h\to 0}\frac{(x+h)^2-x^2}{h}$$

$$= \lim_{h\to 0}(2x+h) = 2x$$

Theorem 6. (Differentiability implies continuity)

Suppose f is differentiable at $x_0 \in D_f$. Then f is continuous at x_0.

Proof. Note that

$$\begin{aligned}
f \text{ is differentiable at } x_0 \in D_f \quad &\Rightarrow \quad \lim_{h\to 0} \tfrac{f(x_0+h)-f(x_0)}{h} = f'(x_0) \\
&\Rightarrow \quad \lim_{h\to 0} f(x_0+h) - f(x_0) = \lim_{h\to 0} h \cdot f'(x_0) \\
&\Rightarrow \quad \lim_{h\to 0} f(x_0+h) - f(x_0) = 0 \\
&\Rightarrow \quad \lim_{h\to 0} f(x_0+h) = f(x_0) \\
&\Rightarrow \quad \lim_{x\to x_0} f(x) = f(x_0) \\
&\Rightarrow \quad f \text{ is continuous at } x_0
\end{aligned}$$

Remark. Every differentiable function is continuous, **but the converse is not true**.

Example 4: Consider the function

$$f(x) = |x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

This function is continuous at all x, but it is not differentiable at $x = 0$.

To see this try to compute the derivative at 0,

- $\lim\limits_{x\to 0^-} \frac{|x|-|0|}{x-0} = \lim\limits_{x\to 0^-} \frac{-x}{x} = -1 \Longrightarrow f_L'(0) = -1.$
- $\lim\limits_{x\to 0^+} \frac{|x|-|0|}{x-0} = \lim\limits_{x\to 0^+} \frac{x}{x} = 1 \Longrightarrow f_R'(0) = 1.$
- note that $f_L'(0) \neq f_R'(0)$.

3.4.2 Differentiability of a function in an interval

Definition.

1. f is said to be differentiable in an open interval $]a, b[$ if it is differentiable at every point x_0 in this interval.

2. f is said to be differentiable in the closed interval $[a, b]$ if

   - f is differentiable in $]a, b[$.
   - f is right differentiable at a point a , i.e. $\lim\limits_{x \to a^+} \frac{f(x)-f(x_0)}{x-x_0} = f^{'}(a)$.
   - f is left continuous at a point b , i.e. $\lim\limits_{x \to b^-} \frac{f(x)-f(x_0)}{x-x_0} = f^{'}(b)$.

3.4.3 Derivative Table of Elementary Functions

$f(x)$	$f^{'}(x)$
a^x	$a^x \ln a$
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
$\tan x$	$\frac{1}{\cos^2 x} = 1 + \tan^2 x$
$\cot x$	$\frac{-1}{\sin^2 x} = -1 - \cot^2 x$
$\arcsin x$	$\frac{1}{\sqrt{1-x^2}}$
$\arccos x$	$\frac{-1}{\sqrt{1-x^2}}$
$\arctan x$	$\frac{1}{1+x^2}$
$\sinh x$	$\cosh x$
$\cosh x$	$\sinh x$

$f(x)$	$f^{'}(x)$
c	0
x	1
$\sqrt{x}$	$\frac{1}{2\sqrt{x}}$
$\sqrt[n]{x}$	$\frac{1}{n.\sqrt[n]{x^{n-1}}}$
$\frac{1}{x}$	$-\frac{1}{x^2}$
$\frac{1}{x^n}$	$\frac{-n}{x^{n+1}}$
x^n	$n.x^{n-1}$
$\ln \lvert x \rvert$	$\frac{1}{x}$
$\exp x$	$\exp x$

3.4.4 Derivative of inverse function

Definition. If f is a function with inverse function f^{-1}, then

$$\left(f^{-1}\right)^{'}(y) = \frac{1}{f^{'}(x)}$$

Example: The inverse of the function $f(x) = x^2$ with reduced domain $[0, +\infty[$ is $f^{-1}(y) = \sqrt{y}$. Use the formula given above to find the derivative of f^{-1}. We have $f^{'}(x) = 2x$, so that $\left(f^{-1}\right)^{'}(y) = \frac{1}{2x} = \frac{1}{2\sqrt{y}}$.

3.4.5 Algebra of derivatives

If f, g are differentiable functions and λ is any constant, then

1. $(f+g)^{'}(x) = f^{'}(x) + g^{'}(x)$.
2. $(f \cdot g)^{'}(x) = f^{'}(x) \cdot g(x) + f(x) \cdot g^{'}(x)$
3. $(\lambda \cdot f)^{'}(x) = \lambda \cdot f^{'}(x)$.
4. $\left(\frac{f}{g}\right)^{'}(x) = \frac{f^{'}(x) \cdot g(x) - f(x) \cdot g^{'}(x)}{(g(x))^2}$, if $g(x) \neq 0$.

3.4.6 Derivatives of composite functions

Definition. If f and g are differentiable, so is the composition $f \circ g$. The derivative of $f \circ g$ is given by:

$$(f \circ g)^{'}(x) = f'(g(x)) \cdot g'(x).$$

Example: The function $f(x) = \sin 2x$ is the composition of two simpler functions, namely: $f(x) = g(h(x))$ where $g(u) = \sin u$ and $h(x) = 2x$. Since g and h are differentiable then $g^{'}(u) = \cos u$ and $h^{'}(x) = 2$.

Therefore the derivative of the composite functions rule implies that

$$f^{'}(x) = (g(h(x)))^{'} = g'(h(x)) \cdot h'(x) = (\cos 2x) \cdot 2 = 2\cos 2x.$$

3.4.7 Derivative of some composite Functions

If f is differentiable function and α is any constant, then:

1. $(f^{\alpha})' = \alpha f' f^{\alpha-1}$, f is strictly positive.
2. $\left(\sqrt{f}\right)' = \frac{f'}{2\sqrt{f}}$, f is strictly positive.
3. $\left(e^{f}\right)' = f' e^{f}$.
4. $(\ln f)' = \frac{f'}{f}$.
5. $(\sin f)' = f' \cos f$.
6. $(\cos f)' = -f' \sin f$.
7. $(\tan f)' = \frac{f'}{\cos^2 f}$.

3.4.8 Lagrange's Mean Value Theorem

Let a, b be two real numbers with $a < b$. Suppose f is a function such that:

(a) f is continuous on $[a, b]$.

(b) f is differentiable on $]a, b[$.

Then there exists $c \in]a, b[$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

3.4.9 L'Hospital's Rule

In this section, we will learn how to evaluate functions whose values cannot be found at certain points.

Indeterminate Forms and L'Hospital's Rule

Consider f and g are differentiable except possibly at x_0. Assume that:



Figure 3.4

(a) $g^{'}(x) \neq 0$ at every point except possibly at x_0.

(b) $\lim\limits_{x \to x_0} \frac{f(x)}{g(x)} = \frac{0}{0}$ or $\lim\limits_{x \to x_0} \frac{f(x)}{g(x)} = \frac{\infty}{\infty}$.

(c) $\lim\limits_{x \to x_0} \frac{f^{'}(x)}{g^{'}(x)}$ exists

$$\text{Then} \quad \lim_{x \to x_0} \frac{f(x)}{g(x)} \quad \text{exists} \quad \text{and} \quad \lim_{x \to x_0} \frac{f^{'}(x)}{g^{'}(x)} = \lim_{x \to x_0} \frac{f(x)}{g(x)}$$

Remark. Note that the rule is also valid for one-sided limits and for limits at infinity or negative infinity.

In fact, for the special case in which $f(x_0) = g(x_0) = 0$, $f^{'}$ and $g^{'}$ are continuous, and $g^{'}(x_0) \neq 0$, it is easy to see why the rule is true.

$$\begin{aligned}
\lim_{x \to x_0} \frac{f^{'}(x)}{g^{'}(x)} &= \frac{f^{'}(x_0)}{g^{'}(x_0)} = \frac{\lim\limits_{x \to x_0} \frac{f(x)-f(x_0)}{x-x_0}}{\lim\limits_{x \to x_0} \frac{g(x)-g(x_0)}{x-x_0}} \\
&= \lim_{x \to x_0} \frac{f(x)-f(x_0)}{g(x)-g(x_0)} \\
&= \lim_{x \to x_0} \frac{f(x)}{g(x)}
\end{aligned}$$

Example 1: Find $\lim\limits_{x \to 1} \frac{\ln x}{x-1}$

$$\lim_{x \to 1} \frac{\ln x}{x-1} = \frac{0}{0}$$

Thus, we can apply l'Hospital's Rule:

$$\lim_{x\to x_0}\frac{f^{'}(x)}{g^{'}(x)} = \lim_{x\to 1}\frac{(\ln x)^{'}}{(x-1)^{'}} = \lim_{x\to 1}\frac{1}{x}$$

$$= 1$$

Then,

$$\lim_{x\to x_0}\frac{f(x)}{g(x)} = \lim_{x\to 1}\frac{\ln x}{x-1} = 1$$

Example 2: Find $\lim\limits_{x\to+\infty}\frac{\exp x}{x^2}$

$$\lim_{x\to+\infty}\frac{\exp x}{x^2} = \frac{\infty}{\infty}$$

Thus, we can apply l'Hospital's Rule:

$$\lim_{x\to x_0}\frac{f^{'}(x)}{g^{'}(x)} = \lim_{x\to+\infty}\frac{(\exp x)^{'}}{(x^2)^{'}} = \lim_{x\to+\infty}\frac{\exp x}{2.x} = \frac{\infty}{\infty}$$

However, a second application of l'Hospital's Rule gives:

$$\lim_{x\to x_0}\frac{f^{''}(x)}{g^{''}(x)} = \lim_{x\to+\infty}\frac{(\exp x)^{'}}{(2.x)^{'}} = \lim_{x\to+\infty}\frac{\exp x}{2} = +\infty$$

Then,

$$\lim_{x\to x_0}\frac{f(x)}{g(x)} = \lim_{x\to+\infty}\frac{\exp x}{x^2} = \infty.$$

Example 3: Find $\lim\limits_{x\to\pi^-}\frac{\sin x}{1-\cos x}$

$$\lim_{x\to\pi^-}\frac{\sin x}{1-\cos x} = \frac{0}{0}$$

If we blindly attempted to use l'Hospital's rule, we would get:

$$\lim_{x\to x_0}\frac{f^{'}(x)}{g^{'}(x)} = \lim_{x\to\pi^-}\frac{(\sin x)^{'}}{(1-\cos x)^{'}} = \lim_{x\to\pi^-}\frac{\cos x}{\sin x} = -\infty$$

Then,

$$\lim_{x\to x_0}\frac{f(x)}{g(x)} = \lim_{x\to\pi^-}\frac{\cos x}{\sin x} = -\infty.$$

3.4.10 Higher derivatives

If the derivative $f^{'}(x)$ of some function f exists for all x in the domain of f, then we have a new function, this function is called the **derivative function** of f, and it is denoted by $f^{'}$. Now that we have agreed that the derivative of a function is a function, we can repeat the process and try to differentiate the derivative. The result, if it exists, is called the **second derivative** of f. It is denoted $f^{''}$. The derivative of the second derivative is called the **third derivative**, written $f^{'''}$, and so on. The n-th derivative of f is denoted $f^{(n)}$. Thus

$$f^{(0)} = f, \quad f^{(1)} = f^{'}, \quad f^{(2)} = \left(f^{'}\right)^{'}, \quad ..., \quad f^{(n)} = \left(f^{n-1}\right)^{'}.$$

Example 1: If $f(x) = x^2 - x + 1$ then

$$\begin{aligned}
f(x) &= x^2 - 2x + 3 \\
f^{'}(x) &= 2x - 2 \\
f^{''}(x) &= 2 \\
f^{'''}(x) &= 0 \\
. &= . \\
. &= . \\
f^{(n)}(x) &= 0.
\end{aligned}$$

Example 2: If $f(x) = \exp x$ then

$$f^{(1)}(x) = \exp x, \quad f^{(2)}(x) = \exp x, ..., \quad f^{(n)} = \exp x.$$

Example 3. If $f(x) = \sin x$ then

$$\begin{aligned}
f^{'}(x) &= \cos x \\
f^{''}(x) &= -\sin x \\
f^{'''}(x) &= -\cos x \\
f^{(4)}(x) &= \sin x \\
f^{(5)}(x) &= \cos x \\
f^{(6)}(x) &= -\sin x \\
f^{(7)}(x) &= -\cos x
\end{aligned}$$

It's easy to find that ,

$$\sin^{(n)} x = \sin\left(x + \frac{n\pi}{2}\right)$$

FAMOUS ISLAMIC MATHEMATICIANS

Sharaf Al-Dīn

Sharaf Al-din dedicated most of his work to **algebra**. He solved extremely bulky **arithmetic** problems using exponents. He founded algebraic identities such as **binomial expansion.** He was the first to define the different degrees of equations and solve them. His contributions have been very useful to the development of Mathematics.

Chapter 4

Finite Expansions

4.1 Approximating functions by Taylor polynomials

4.1.1 First degree Taylor polynomials

If we know the function value at some point $f(x_0)$ and the value of the derivative at the same point $f^{'}(x_0)$, we can use these to find the tangent line, and then use the tangent line to approximate $f(x)$ for other points x.

The tangent line approximation of f for x near x_0 is called **the first degree Taylor polynomial** of f and is:

$$f(x) \approx f(x_0) + f^{'}(x_0)\,(x - x_0)\,. \qquad (\blacklozenge)$$

The statement that a complicated function behaves like a simpler function f for x near x_0 can be made more precise by use of the "O" notation. For example, we can replace the weak statement ($\blacklozenge$) by the stronger version,

$$f(x) = f(x_0) + f^{'}(x_0)\,(x - x_0) + O(x - x_0)$$

This means that there exists a function $\varepsilon(x)$ such that:

$$\lim_{x \to x_0} \frac{(x - x_0).\varepsilon(x)}{(x - x_0)} = 0$$

we write then

$$(x - x_0).\varepsilon(x) = O(x - x_0)$$

Example 1: Consider the function $f(x) = \sin x$. We want the first degree Taylor polynomial of this function near the point $x_0 = \frac{\pi}{4}$ and $x_0 = 0$.

- Since $\sin(\frac{\pi}{4}) = \frac{\sqrt{2}}{2}$ and $(\sin)'(\frac{\pi}{4}) = \cos(\frac{\pi}{4}) = \frac{\sqrt{2}}{2}$, the approximation of f for x near $x_0 = \frac{\pi}{4}$ is given by:

$$\sin x = \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}\left(x - \frac{\pi}{4}\right) + O(x - \frac{\pi}{4}), \quad \lim_{x \to \frac{\pi}{4}} O(x - \frac{\pi}{4}) = 0$$

- Since $\sin(0) = 1$ and $(\sin)'(0) = \cos 0 = 1$, the approximation of f for x near $x_0 = 0$ is given by:

$$\sin x = x + O(x), \quad \lim_{x \to 0} O(x) = 0$$

Example 2: Consider the function $f(x) = \sqrt{x}$. We want the first degree Taylor polynomial of this function near the point $x_0 = 1$ and an evaluation of $\sqrt{1.002}$

the approximation of f for x near $x_0 = 1$ is given by:

$$\sqrt{x} = 1 + \frac{1}{2}(x - 1) + O(x - 1), \quad \lim_{x \to 0} O(x - 1) = 0$$

then

$$\sqrt{1.002} = 1 + \frac{1}{2}(1.002 - 1) + O(0.002) = 1.001 + O(0.002), \quad \lim_{x \to 0} O(0.002) = 0.$$

Higher order Taylor polynomials

The approximations of the function f by **the Taylor polynomial of degree** n, denoted by $P_n(x - x_0)$ for x near x_0 using more derivatives $f'(x_0), f''(x_0), ..., f^{(n)}(x_0)$ is given by: $f(x) = P_n(x - x_0) + R_n(x - x_0)$ for $\lim_{x \to x_0} R_n(x - x_0) = 0$.

Note that $R_n(x - x_0)$ is called **remainder term** which is the **approximation error** when approximating f

with its Taylor polynomial. Using the "O" notation, the statement in Taylor polynomial reads as

$$R_n\left(x-x_0\right)=O\left(\left(x-x_0\right)^n\right)$$

$$\boxed{f(x)=f(x_0)+\frac{f^{'}(x_0)}{1!}\left(x-x_0\right)+\frac{f^{''}(x_0)}{2!}\left(x-x_0\right)^2+...+\frac{f^{(n)}(x_0)}{n!}\left(x-x_0\right)^n+O\left(\left(x-x_0\right)^n\right)}$$

Example 1:

Consider the function $f(x)=\ln x$. We want a polynomial approximation of this function near the point $x_0=1$.

The first few derivatives of f are

$$f(x)=\ln x$$

$$f^{'}(x)=\tfrac{1}{x}$$

$$f^{''}(x)=\tfrac{-1}{x^2}$$

$$f^{'''}(x)=\tfrac{2}{x^3}$$

$$f^{(4)}(x)=\tfrac{-3}{x^4}$$

The derivatives evaluated at $x_0=1$ are

$$f(0)=0, \qquad f^{(n)}(0)=(-1)^{n-1}(n-1)!$$

By Taylor's polynomial we have,

$$\ln x=(x-1)-\frac{(x-1)^2}{2}+\frac{(x-1)^3}{3}-\frac{(x-1)^4}{4}+...+(-1)^{n-1}\frac{(x-1)^n}{n}+O\left((x-1)^n\right)$$

4.1.2 Mac-Laurin Polynomials

The approximations of f by the Taylor polynomial of degree n for x near $x_0 = 0$ is called the approximations of f by **Mac-Laurin polynomial** and is given by:

$$\boxed{f(x) = f(0) + \frac{f^{'}(0)}{1!}x + \frac{f^{''}(0)}{2!}x^2 + ... + \frac{f^{(n)}(0)}{n!}.x^n + O\left(x^n\right), \lim_{x\to 0} O\left(x^n\right) = 0.}$$

Example: Consider the function $f(x) = \cos x$. We want Mac-Laurin polynomial of this function near the point $x_0 = 0$. The first few derivatives of f are

$$\begin{aligned}
f^{'}(x) &= \cos x \\
f^{''}(x) &= -\sin x \\
f^{'''}(x) &= -\cos x \\
f^{(4)}(x) &= \sin x \\
f^{(5)}(x) &= \cos x \\
f^{(6)}(x) &= -\sin x
\end{aligned}$$

It's easy to find that ,

$$\cos^n(x) = \cos\left(x + \frac{n\pi}{2}\right)$$

Since $\cos(0) = 1$ and $\sin(0) = 0$ the Maclaurin polynomials of the cosine is,

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + ... + (-1)^{2n}\frac{x^{2n}}{2n!} + O\left(x^{2n+1}\right)$$

The polynomial approximation of degree 4 is given by:

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + O\left(x^4\right)$$

The polynomial approximation of degree 5 is given by:

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + O\left(x^5\right)$$

The first statement informs us that there are terms of order x^4 in the expansion. The second statement is stronger as it informs us that there are no terms of order x^5.

4.2 Finite expansions at zero

Definition.

Let f be a real valued function. We said that the function f is represented by **a finite expansion at zero** if there exist real numbers $a_{0,}a_{1,}...,a_n$ and a real valued function ε such that

$$\boxed{f(x) = a_0 + a_1 x + a_2 x^2 + ... + a_n x^n + x^n \varepsilon(x),\ \lim_{x \to 0} \varepsilon(x) = 0.}$$

Then the function f is represented by the polynomial approximation of degree n, denoted by $P_n(x)$ for x near zero, which is called *the main part* of finite expansions at zero, such that: $P_n(x) = a_0 + a_1 x + a_2 x^2 + ... + a_n x^n$.

Remark: Note that $x^n \varepsilon(x) = O\left(x^n\right)$.

Example. Using the euclidean division by increasing power order, one has the finite expansion at zero of $f(x) = \frac{1}{1-x}$:

$$\frac{1}{1-x} = 1 + x + x^2 + ... + x^n + \frac{x^{n+1}}{1-x} = 1 + x + x^2 + ... + x^n + x^n . \left(\frac{x}{1-x}\right).$$

in this case $\varepsilon(x) = \frac{x}{1-x}$. We generally do not try to determine the function $\varepsilon(x)$.

Properties.

(a) If the function f can be expanded at zero, then this expansion is *unique*.

(b) If the function f can be expanded at zero, then $\lim\limits_{x\to 0} f(x)$ exists and equal to a_0. This criterion is generally used to demonstrate that a function does not admit an expansion.

Example. The function $f(x) = \ln x$ does not have an expansion at zero, because $\lim\limits_{x\to 0^+} f(x) = -\infty$.

4.2.1 Basic finite expansions of elementary functions

$\exp x$	$= 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + ... + \frac{x^n}{n!} + O\left(x^n\right).$
$(1+x)^\alpha$	$= 1 + \alpha x + \frac{\alpha(\alpha-1)x^2}{2!} + ... + \frac{\alpha(\alpha-1)....(\alpha-(n-1))x^n}{n!} + O\left(x^n\right).$
$\frac{1}{1+x}$	$= 1 - x + x^2 - x^3 + x^4 + ... + (-1)^n x^n + O\left(x^n\right).$
$\frac{1}{1-x}$	$= 1 + x + x^2 + x^3 + x^4 + ... + x^n + O\left(x^n\right).$
$\cos x$	$= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + ... + \frac{(-1)^n x^{2n}}{2n!} + O\left(x^{2n+1}\right).$
$\sin x$	$= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + ... + \frac{(-1)^n x^{2n+1}}{2n+1!} + O\left(x^{2n+1}\right).$
$\ln(1+x)$	$= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + ... + \frac{(-1)^{n+1} x^n}{n} + O\left(x^n\right).$
$\ln(1-x)$	$= -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - .... - \frac{x^n}{n} + O\left(x^n\right).$
$\arccos x$	$= \frac{\pi}{2} - x - \frac{1}{2}.\frac{x^3}{3.} - \frac{1.3}{2.4}.\frac{x^5}{5} - \frac{1.3.5}{2.4.6}\frac{x^7}{7} - ... - \frac{1.3.5...(2n-1).}{2.4.6....2n}.\frac{x^{(2n+1)}}{(2n+1)} + O\left(x^{2n+1}\right).$
$\arcsin x$	$= x + \frac{1}{2}.\frac{x^3}{3.} + \frac{1.3}{2.4}.\frac{x^5}{5} + \frac{1.3.5}{2.4.6}\frac{x^7}{7} + ... + \frac{1.3.5...(2n-1).}{2.4.6....2n}.\frac{x^{(2n+1)}}{(2n+1)} + O\left(x^{2n+1}\right).$
$\arctan g$	$= x - \frac{x^3}{3.} - \frac{x^5}{5} - \frac{x^7}{7} + ... - \frac{x^{(2n+1)}}{(2n+1)} + O\left(x^{2n+1}\right).$
$\cosh x$	$= 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + ... + \frac{x^{2n}}{2n!} + O\left(x^{2n}\right).$
$\sinh x$	$= x + \frac{x^3}{3!} + \frac{x^5}{5!} + ... + \frac{x^{2n+1}}{2n+1!} + O\left(x^{2n+1}\right).$

4.3 Algebraic combinations of finite expansions

Definition.

If f and g can both be expanded at zero and λ is any constant, then each of the following functions is also can be expanded at zero: The sum $f+g$, the difference $f-g$, the constant multiple λf , the product $f \cdot g$, the quotient f/g , if $g(x_0) \neq 0$.

Consider the finite expansions at zero of f and g:

$$f(x) = a_0 + a_1 x + a_2 x^2 + ... + a_n x^n + O\left(x^n\right)$$

$$g(x) = b_0 + b_1 x + b_2 x^2 + ... + b_n x^n + O\left(x^n\right)$$

★ The finite expansion at zero of the sum $f+g$ is:

$$\begin{aligned} (f+g)(x) &= (a_0+b_0) + (a_1+b_1)x + (a_2+b_2)x^2 + ... + (a_n+b_n)x^n + O\left(x^n\right) \\ \lim_{x\to 0} \varepsilon(x) &= 0. \end{aligned}$$

★ The finite expansion at zero of the $f \cdot g$ is obtained by the product and keeping only the monomials of degree less than n in the product

$$\left(a_0 + a_1 x + a_2 x^2 + ... + a_n x^n\right)\left(b_0 + b_1 x + b_2 x^2 + ... + b_n x^n\right)$$

★ The finite expansion at zero of the quotient f/g is obtained by the euclidean division of $\left(a_0 + a_1 x + a_2 x^2 + ... + a_n x^n\right)$ by $\left(b_0 + b_1 x + b_2 x^2 + ... + b_n x^n\right)$ by increasing power order.

Example 1: Find the finite expansion at zero of $f(x) = \sinh x$ of the degree 4.

$$\begin{aligned}
\sinh x &= \frac{\exp x - \exp(-x)}{2} \\
&= \frac{1}{2}\left[\left(1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!}\right) - \left(1 + \frac{-x}{1!} + \frac{x^2}{2!} + \frac{-x^3}{3!} + \frac{x^4}{4!}\right)\right] + O\left(x^4\right) \\
&= \frac{1}{2}\left(2x + 2\frac{x^3}{3!}\right) + O\left(x^4\right) \\
&= x + \frac{x^3}{3!} + O\left(x^4\right).
\end{aligned}$$

Example 2: Find the finite expansion at zero of $f(x) = \cos x \cdot \sin x$ of the degree 5.

We have $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + O\left(x^5\right)$ and $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} + O\left(x^5\right)$.

$$\begin{aligned}
f(x) &= \cos x \cdot \sin x \\
&= \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!}\right) \cdot \left(x - \frac{x^3}{3!} + \frac{x^5}{5!}\right) + O\left(x^5\right). \\
&= \left(1 - \frac{x^2}{2} + \frac{x^4}{24}\right) \cdot \left(x - \frac{x^3}{6} + \frac{x^5}{120}\right) + O\left(x^5\right). \\
&= x - \frac{2}{3}x^3 + \frac{2}{15}x^5 + O\left(x^5\right).
\end{aligned}$$

Example 3: Find the finite expansion at zero of $f(x) = \frac{\sin x}{\cos x}$ of the degree 3.

Note that $\lim\limits_{x \to 0} \cos x \neq 0$ then the quotient $f(x) = \frac{\sin x}{\cos x}$ can be expanded at zero.

Let $\sin x = x - \frac{x^3}{3!} + O\left(x^3\right)$ and $\cos x = 1 - \frac{x^2}{2!} + O\left(x^3\right)$.

Using the euclidean division by increasing power order we obtain:

$$\begin{aligned}
f(x) &= \frac{\sin x}{\cos x} \\
&= \frac{x - \frac{x^3}{3!}}{1 - \frac{x^2}{2!}} + O\left(x^3\right) \\
&= x + \frac{1}{3}x^3 + O\left(x^3\right).
\end{aligned}$$

Example 4: Find the finite expansion at zero of $f(x) = \frac{\ln(1+x)}{\sin x}$ of the degree 3.

Since $\sin x = x - \frac{x^3}{3!} + O\left(x^3\right)$ we have $\lim\limits_{x \to 0} \sin x = 0$.

Note that the function f can be expanded at zero of the degree 3 if the finite expansions at zero of $\ln(1+x)$ and $\sin x$ are given of the degree 4.

$$\begin{aligned} f(x) &= \frac{\ln(1+x)}{\sin x} \\ &= \frac{x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + O\left(x^4\right)}{x - \frac{x^3}{3!} + O\left(x^4\right)} \\ &= \frac{1 - \frac{x}{2} + \frac{x^2}{3} - \frac{x^3}{4} + O\left(x^3\right)}{1 - \frac{x^2}{3!} + O\left(x^3\right)} \end{aligned}$$

Since $\lim\limits_{x\to 0}\left(1 - \frac{x^2}{3!} + O\left(x^3\right)\right) = 0$ the function f can be expanded in this case.

$$\begin{aligned} \frac{\ln(1+x)}{\sin x} &= \frac{1 - \frac{x}{2} + \frac{x^2}{3} - \frac{x^3}{4} + O\left(x^3\right)}{1 - \frac{x^2}{3!} + O\left(x^3\right)} \\ &= 1 - \frac{x}{2} + \frac{x^2}{6} - \frac{x^3}{12} + O\left(x^3\right). \end{aligned}$$

4.4 Composite of finite expansions

Definition.

If g can be expanded at zero of degree n and if f can be expanded at $g(0)$ of degree n such that $g(0) = 0$. Then the composite function $(f \circ g)$ can be expanded at zero of degree n by replacing the finite expansion of g in the finite expansion of f and by keeping only the monomials of degree $\leq n$.

Example 1: Find the finite expansion at zero of $f(x) = \exp(\cos x)$ of the degree 3.

If $g(x) = \cos x$ note that $g(0) \neq 1$.

We know that $\exp x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + O\left(x^3\right)$ and $\cos x = 1 - \frac{x^2}{2!} + O\left(x^3\right)$. So if $g(x) = \cos x - 1 =$

$-\frac{x^2}{2!} + O\left(x^3\right)$ in this case $g(0) = 0$.

$$\begin{aligned}
f(x) &= \exp(\cos x) \\
&= \exp(1 - 1 + \cos x) \\
&= \exp 1 \cdot \exp(-1 + \cos x) \\
&= \exp 1 \cdot \exp\left[-\frac{x^2}{2!} + O\left(x^3\right)\right] \\
&= \exp 1. \left[1 + \frac{\left(-\frac{x^2}{2!}\right)}{1!} + O\left(x^3\right)\right] \\
&= \exp 1 - \frac{\exp 1}{2} x^2 + O\left(x^3\right).
\end{aligned}$$

Example 2: Prove that the finite expansion at zero of $f(x) = \exp(\sin x)$ is given by $f(x) = \exp(\sin x) = 1 + x + \frac{x^2}{2} + O\left(x^3\right)$.

4.5 Finite expansions at a point

We said that the function $f : x \longmapsto f(x)$ can be represented by a finite expension at point x_0 if the function $F : X \longmapsto F(X)$ can be represented by finite expension at zero $X_0 = 0$ such that $F(X) = f(x_0 + X)$ and

$$F(X) = a_0 + a_1 X + a_2 X^2 + ... + a_n X^n + O(X^n), \lim_{X \to 0} O(X^n) = 0.$$

$$f(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)^2 + ... + a_n(x - x_0)^n + O((x - x_0)^n),$$

and $\lim_{x \to x_0} O((x - x_0)^n) = 0$.

Example 1: Find the finite expansion at a point $x_0 = 1$ of $f(x) = \exp x$ of the degree 3 .

$$\begin{aligned}
F(X) &= f(x_0 + X) \\
&= \exp(1 + X) \\
&= \exp 1 \cdot \exp X \\
&= \exp 1 \cdot \left[1 + \frac{X}{1!} + \frac{X^2}{2!} + \frac{X^3}{3!} + O\left(X^3\right)\right] \\
&= \exp 1 \cdot \left[1 + \frac{(x-1)}{1!} + \frac{(x-1)^2}{2!} + \frac{(x-1)^3}{3!} + O\left((x-1)^3\right)\right]
\end{aligned}$$

Example 2: Find the finite expansion at a point $x_0 = 2$ of $f(x) = \ln x$ of the degree 2 .

$$\begin{aligned}
F(X) &= f(x_0 + X) \\
&= \ln(2 + X) \\
&= \ln\left[2 \cdot \left(1 + \frac{X}{2}\right)\right] \\
&= \ln 2 + \ln\left(1 + \frac{X}{2}\right) \\
&= \ln 2 + \frac{1}{2}X - \frac{1}{8}X^2 + O\left(X^2\right) \\
&= \ln 2 + \frac{1}{2}(x-2) - \frac{1}{8}(x-2)^2 + O\left((x-2)^2\right).
\end{aligned}$$

4.6 Finite expansions at Infinity

We said that the function $f : x \longmapsto f(x)$ can be represented by a finite expension at infinity if the function $F : X \longmapsto F(X)$ can be represented by finite expension at zero $X_0 = 0$ such that $F(X) = f\left(\frac{1}{x}\right)$ and

$$F(X) = a_0 + a_1 X + a_2 X^2 + ... + a_n X^n + O(X^n), \lim_{X \to \infty} O(X^n) = 0.$$

$$f(x) = a_0 + \frac{a_1}{x} + \frac{a_2}{x^2} + ... + \frac{a_n}{x^n} + O\left(\frac{1}{x^n}\right).$$

Example 1: Find the finite expansion at infinity of $f(x) = \cos \frac{1}{x}$.

Let $X = \frac{1}{x}$ and thus:

$$\begin{aligned}\cos\frac{1}{x} &= \cos X = 1 - \frac{X^2}{2!} + \frac{X^4}{4!} + ... + \frac{(-1)^n X^{2n}}{2n!} + O\left(X^{2n}\right) \\ &= 1 - \frac{1}{2!x^2} + \frac{1}{4!x^4} + ... + \frac{(-1)^n}{2n!x^{2n}} + O\left(\frac{1}{x^{2n}}\right)\end{aligned}$$

4.7 Using finite expansions to evaluate limits

The finite expansions provide a good way to understand the behaviour of a function near a specified point and so are useful for solving some indeterminate forms. When taking a limit as $x \to 0$, we can often simplify the statement by substituting in finite expansions that we know.

Example 1: Find the limit $\lim\limits_{x\to 0} \frac{\exp(2x)\sin 3x}{\sinh(-2x)}$

$$\lim_{x\to 0} \frac{\exp(2x)\sin 3x}{\sinh(-2x)} = \lim_{x\to 0} \frac{(1+2x)\cdot(3x) + O(x)}{-2x + O(x)} = -\frac{3}{2}.$$

Example 2: Find the limit $\lim\limits_{x\to 0} \frac{1-\cos x}{\sin^2 x}$

$$\lim_{x\to 0} \frac{1-\cos x}{\sin^2 x} = \lim_{x\to 0} \frac{\frac{x^2}{2} + O(x)}{(x+O(x))^2} = \frac{1}{2}.$$

Example 3: Find the limit $\lim\limits_{x\to 0} \frac{\exp x - 1 - x - \frac{x^2}{2}}{x^3}$

$$\exp x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + O\left(x^3\right)$$

We get $\exp x - 1 - x - \frac{x^2}{2} = \frac{x^3}{3!} + O\left(x^3\right)$

and consequently

$$\begin{aligned}\frac{\exp x - 1 - x - \frac{x^2}{2}}{x^3} &= \frac{\frac{x^3}{3!} + O\left(x^3\right)}{x^3} \\ &= \frac{1}{3!} + O\left(x\right)\end{aligned}$$

so

$$\lim_{x\to 0}\frac{\exp x-1-x-\frac{x^2}{2}}{x^3}=\lim_{x\to 0}\frac{1}{3!}+O\left(x\right)=\frac{1}{6}.$$

FAMOUS ISLAMIC MATHEMATICIANS



Abu Reyhan Al-Biruni

Al-Biruni is considered one of Islamic mathematical genius. He discovered **mathematical techniques** to determine exactly the beginnings of the season. He also wrote about the sun and its movements and the eclipse. He combined **trigonometry** and **algebra** to achieve this very numerical feat.

Chapter 5

Vector Space and Linear Maps

5.1 Vector Space

Underlying every **vector space** (to be defined shortly) is **a scalar field** K.

A **field** is a set of elements where the four basic operations $+, -, \times, \div$ are defined, with their usual properties (commutativity, associativity, distributivity). Examples of fields include the rational numbers $\mathbb{Q}$, the real numbers $\mathbb{R}$, and the complex numbers $\mathbb{C}$. However, $\mathbb{N}$ is not a field (we cannot subtract or divide) and $\mathbb{Z}$ is not a field (we cannot divide).

Definition. (**Vector space**)

A **vector space** over a field K is a nonempty set V of objects, called vectors, on which are defined two operations:

1) *An internal operation* (*vector addition)*

$$\begin{aligned} + : \quad V \times V &\longrightarrow \quad V \\ (u, v) &\longmapsto \quad u + v \end{aligned}$$

2) *An external operation* (*scalar multiplication)*

$$\begin{array}{rccc} \cdot : & K \times V & \longrightarrow & V \\ & (\alpha, u) & \longmapsto & \alpha u \end{array}$$

such that the following properties are satisfied:

(a) $\forall u, v \in V, u + v = v + u.$

(b) $\forall u, v, w \in V : (u + v) + w = u + (v + w)\,.$

(c) $\forall u \in V$: $u + 0_V = u$ (0_V is the zero vector).

(d) $\forall u \in V, \exists\,(-u) \in V :\ u + (-u) = 0_V.$

(e) $\forall u, v \in V, \forall \alpha \in K : \alpha\,(u + v) = \alpha u + \alpha v.$

(f) $\forall u \in V, \forall \alpha, \beta \in K : (\alpha + \beta)\,u = \alpha u + \beta u.$

(g) $\forall u \in V, \forall \alpha, \beta \in K : (\alpha\beta)\,u = \alpha\,(\beta u)\,.$

(h) $\forall u \in V : 1u = u.$

Examples:

(a) $K^n = \{(x_1, x_2, .., x_n)\,/x_i \in \mathbb{R}, i = 1, .., n\}$ is a vector space over the field K and K is any field (typically $K = \mathbb{R}$ or $K = \mathbb{C}$) with the vector addition and scalar multiplication defined as follows for all $(x_1, x_2, .., x_n)$ and $(y_1, y_2, .., y_n)$ from K^n and $\alpha \in K$:

$$\begin{aligned} (x_1, x_2, .., x_n) + (y_1, y_2, .., y_n) &= (x_{1+}y_1, x_{2+}y_2, .., x_n + y_n) \\ \alpha\,(x_1, x_2, .., x_n) &= (\alpha x_1, \alpha x_2, .., \alpha x_n) \end{aligned}$$

(b) The set $P[x] = \left\{\sum_{i=1}^{n} a_i x^i /\ a_i \in \mathbb{R}, i = 1, .., n\right\}$ of all polynomials over a field $\mathbb{R}$ is a vector space over $\mathbb{R}$ with the vector addition and scalar multiplication defined as follows for all $p(x) = \sum_{i=1}^{n} a_i x^i$ and $q(x) = \sum_{i=1}^{n} b_i x^i$ from $P[x]$ and $\alpha \in \mathbb{R}$:

$$\begin{aligned}\forall p(x), q(x) &\in P[x] : p(x) + q(x) = \sum_{i=1}^{n} (a_i + b_i)\, x^i \\ \forall p(x) &\in P[x], \forall \alpha \in \mathbb{R} : \alpha p(x) = \sum_{i=1}^{n} (\alpha a_n)\, x^n\end{aligned}$$

(c) The set V of all real valued continuous (differentiable or integrable) functions defined on the closed interval $[a, b]$ is a real vector space with the vector addition and scalar multiplication defined as follows:

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\ (\alpha f)(x) &= \alpha f(x)\end{aligned}$$

For all $f, g \in V$ and $\alpha \in \mathbb{R}$.

5.1.1 Subspaces of a vector space

Definition 1. (Subspace)

A subspace of a vector space V is a nonempty subset F of V that has two properties:

(a) F is closed under vector addition. That is, $\forall u, v \in F : u + v \in F$.

(b) F is closed under multiplication by scalars. That is, $\forall u \in F, \forall \alpha \in K : \alpha u \in F$.

Definition 2. (Subspace)

A subspace of a vector space V is a a nonempty subset F of V if and only if:

$$\forall u, v \in F, \forall \alpha, \beta \in K : \alpha u + \beta v \in F.$$

Remark 1.

Properties (1) and (2) guarantee that a subspace F of V is itself a vector space, under the vector space operations already defined in V.

Example 1:

- $\mathbb{R}^{n-1}$ is a subspace of $\mathbb{R}^n$.
- $\{0_V\}$ is a subspace of V.
- V is a subspace of V.

Example 2: Show that $F = \{(0, y, z), y, z \in \mathbb{R}\}$ is a subspace of real vector space $\mathbb{R}^3$.

- $0_{\mathbb{R}^3} \in F$ then F is a nonempty subset of $\mathbb{R}^3$.
- Let $u = (0, y_1, z_1)$, $v = (0, y_2, z_2) \in F$ and $\alpha, \beta \in \mathbb{R}$. Then,

$$\begin{aligned} \alpha u + \beta v &= \alpha (0, y_1, z_1) + \beta (0, y_2, z_2) \\ &= (0, \alpha y_1 + \beta y_2, \alpha z_1 + \beta z_2) \in F. \end{aligned}$$

Hence, F is a subspace of $\mathbb{R}^3$.

5.1.2 Linear combinations

Definition. (Linear combination)

Let V be a vector space. We say that the vector u is a **linear combination** of the vectors $v_1, v_2, .., v_n$ of V if

$$\exists\, \alpha_1, \alpha_2, .., \alpha_n \in K : u = \alpha_1 v_1 + \alpha_2 v_2 + .. + \alpha_n v_n.$$

Example: Express $u = (-2, 3)$ in $\mathbb{R}^2$ over $\mathbb{R}$ as a linear combination of the vectors $v_1 = (1, 1)$ and $v_2 = (1, 2)$.

Let α_1, α_2 be scalars such that

$$
\begin{array}{rcl}
u & = & \alpha_1 v_1 + \alpha_2 v_2 \\
\Rightarrow \quad (-2,3) & = & \alpha_1(1,1) + \alpha_2(1,2) \\
\Rightarrow \quad (-2,3) & = & (\alpha_1 + \alpha_2, \alpha_1 + 2\alpha_2) \\
\Rightarrow \quad \alpha_1 + \alpha_2 = -2 & \text{and} & \alpha_1 + 2\alpha_2 = 3 \\
\Rightarrow \quad \alpha_1 = -7 & \text{and} & \alpha_2 = 5
\end{array}
$$

Hence, $u = -7v_1 + 5v_2$

5.1.3 Linear independence and linear dependence

Definition. (**Linear independence**)

Let the set $S = \{v_1, v_2, .., v_n\} \subset V$, a vector space. We say that S is **linearly independent** if all scalars $\alpha_1, \alpha_2, .., \alpha_n$ are zero for which $\alpha_1 v_1 + \alpha_2 v_2 + .. + \alpha_n v_n = 0_V$. That is

$$\forall \alpha_1, \alpha_2, .., \alpha_n \in K : \alpha_1 v_1 + \alpha_2 v_2 + .. + \alpha_n v_n = 0_V \Rightarrow \alpha_1 = \alpha_2 = .. = \alpha_n = 0_K.$$

Otherwise we say S is **linearly dependent.**

Definition. (**Linear dependence**)

The set $S = \{v_1, v_2, .., v_n\}$ is **linearly dependent** if there are scalars $\alpha_1, \alpha_2, .., \alpha_n$ not all zero for which

$$\alpha_1 v_1 + \alpha_2 v_2 + .. + \alpha_n v_n = 0_V$$

That is:

$$\exists\, \alpha_1, \alpha_2, .., \alpha_n \in K : \alpha_i \neq 0, i \in \{1, .., n\} \wedge \alpha_1 v_1 + \alpha_2 v_2 + .. + \alpha_n v_n = 0_V.$$

Example 1: The set $S = \{(-1,0), (2,1)\}$ is **linearly independent.**

Let $\alpha_1, \alpha_2 \in \mathbb{R}$:

$$\begin{aligned} \alpha_1 (-1,0) + \alpha_2 (2,1) = (0,0) \quad &\Rightarrow \quad (-\alpha_1 + 2\alpha_2, \alpha_2) = (0,0)\,. \\ &\Rightarrow \quad -\alpha_1 + 2\alpha_2 = 0 \text{ and } \alpha_2 = 0. \\ &\Rightarrow \quad \alpha_1 = \alpha_2 = 0. \end{aligned}$$

Example 2: The set $S = \{(1,0), (-2,0)\}$ is **linearly dependent.**

Let $\alpha_1, \alpha_2 \in \mathbb{R}$:

$$\begin{aligned} \alpha_1 (1,0) + \alpha_2 (-2,0) = (0,0) \quad &\Rightarrow \quad (\alpha_1 - 2\alpha_2, 0) = (0,0)\,. \\ &\Rightarrow \quad \alpha_1 - 2\alpha_2 = 0 \\ &\Rightarrow \quad \alpha_1 = 2\alpha_2. \end{aligned}$$

$$\exists \alpha_1 = 1 \wedge \alpha_2 = \tfrac{1}{2} \qquad \wedge \quad \alpha_1 (1,0) + \alpha_2 (-2,0) = (0,0)\,.$$

Remark 2.

- $\{u\}$ is **linearly independent** $\Leftrightarrow u \neq 0_V$.
- $0_V \in S = \{v_1, v_2, .., v_n\} \Longrightarrow S$ is **linearly dependent.**

5.1.4 Generating sets

Definition. **(Generating sets)**

Given a vector space V, a finite set of vectors $S = \{v_1, v_2, .., v_n\} \subset V$ is called a system of **generators** if every vector $u \in V$ can be expressed as a linear combination of vectors of S :

$$\forall u \in V, \exists\, \alpha_1, \alpha_2, .., \alpha_n \in K : u = \alpha_1 v_1 + \alpha_2 v_2 + .. + \alpha_n v_n.$$

and we write $V = [S]$.

Example 1: The set $S = \{(1,1,1), (2,2,0), (3,0,0)\}$ is a system of generators of $\mathbb{R}^3$.

Let $u = (x, y, z)$ be a vector, we check the scalars $\alpha, \beta, \gamma \in \mathbb{R}$ such that

$$u = (x, y, z) = \alpha (1, 1, 1) + \beta (2, 2, 0) + \gamma (3, 0, 0)$$

$$\Rightarrow \begin{cases} x = \alpha + 2\beta + 3\gamma \\ y = \alpha + 2\beta \\ z = \alpha \end{cases} \Rightarrow \begin{cases} \alpha = z \\ \beta = \frac{y-z}{2} \\ \gamma = \frac{x-y}{3} \end{cases}$$

$$\Rightarrow \quad u = z (1, 1, 1) + \frac{y - z}{2} (2, 2, 0) + \frac{x - y}{3} (3, 0, 0)$$

Example 2: Find generating set of the vector space $\mathbb{R}^2$ over the field $\mathbb{R}$

$$(x, y) = x (1, 0) + y (0, 1) \Rightarrow \mathbb{R}^2 = [\{(1, 0), (0, 1)\}]$$

5.1.5 Bases of a vector space

Definition. **(Bases)**

Given a vector space V, a finite set of vectors $S = \{v_1, v_2, .., v_n\} \subset V$ is called a **basis** of V if has two properties:

(a) S is **linearly independent.**

(b) S is a **generator of** V, that is $V = [S]$.

Examples: Let $\mathbb{R}^n$ be a vector space over $\mathbb{R}$.

- If $n = 1$, the basis of $\mathbb{R}$ is the set $S_1 = \{1\}$.
  - Since $\forall x \in \mathbb{R} : x = 1.x$ then $\mathbb{R}$ is generated by S_1.
  - $\forall \alpha \in \mathbb{R} : \alpha.1 = 0 \Rightarrow \alpha = 0$. Thus the set S_1 is linearly independent.

- If $n = 2$, the basis of $\mathbb{R}^2$ is the set $S_2 = \{(1,0), (0,1)\}$.
  - $\mathbb{R}^2$ is generated by S_2.
  - $\forall \alpha, \beta \in \mathbb{R} : \alpha.(1,0) + \beta.(0,1) = (\alpha, \beta) = (0,0) \Rightarrow \alpha = \beta = 0$. Thus the set S_2 is linearly independent.
- If $n = 3$, the basis of $\mathbb{R}^3$ is the set $S_3 = \{(1,0,0), (0,1,0), (0,0,1)\}$.
  - Clearly S_3 is linearly independent and is a system of generators of $\mathbb{R}^3$.

Property. If S is a basis, every vector can be written as a linear combination of its elements in a *unique way*.

Example 1. Let the set $S = \{(1,0), (1,1)\}$, we can write any vector of $\mathbb{R}^2$ as a linear combination of $(1,0)$ and $(1,1)$ in a unique way.

Example 2. The set $S = \{(1,0), (0,1), (1,1)\}$ is not a basis but it is a system of generators. In this case any of the three vectors can be removed because it can be expressed as a combination of the other two. The linear combinations are not unique:

$$(2,3) = 1.(1,0) + 2.(0,1) + 1.(1,1)$$

5.1.6 Dimension of a vector space

Definition. (**Dimension**)

If a vector space V has a basis with finite number of elements, then every other basis of V has the same number of elements. This number is called the dimension of V. We write

$$\dim V = n$$

Properties. if we know that a vector space has dimension n $(\dim V = n)$, then:

- Every basis consists of exactly n vectors (but not every set of n vectors is a basis!).
- Every system of generators has to contain at least n vectors.
- If a system of generators consists of n vectors, then it is a basis.

- If a set of n vectors is linearly independent, then it is a basis.

Proposition. Let V be a vector space of dimension n, and let F is a subspace of a vector space V then $\dim F \leq \dim V$. Furthermore, if $\dim F = \dim V$ then $F = V$.

Example 1: $\mathbb{R}^n$ is a vector space of dimension n. A particular basis is the *canonical basis*:

$$\begin{aligned} e_1 &= (1, 0, 0, .., 0) \\ e_2 &= (0, 1, 0, .., 0) \\ &. \\ &. \\ e_n &= (0, 0, 0, .., 1) \end{aligned}$$

Example 2: The set of three vectors $\{(1, 2), (-1, 2), (3, 1)\}$ is not a basis of $\mathbb{R}^2$ because $3 > \dim \mathbb{R}^2 = 2$**.**

Example 3:

- $\dim \{0_V\} = 0$.
- Let $\mathbb{C}$ be a vector space over $\mathbb{R}$, then $\dim \mathbb{C} = 2$.
- Let P_n be a vector space of polynomial over $\mathbb{R}$, then $\dim(P_n) = n + 1$.

5.2 Linear Maps

Definition. (**Linear map**)

A **linear map** f from a vector space V into a vector space W over the same field K is a rule that assigns to each vector x in V a unique vector $f(x)$ in W, such that:

$$\begin{aligned} f: \ V &\longrightarrow W \\ x &\longmapsto f(x) \end{aligned}$$

(a) $\forall x_1, x_2 \in V : f(x_1 + x_2) = f(x_1) + f(x_2)$.

(b) $\forall x \in V, \forall \alpha \in K : f(\alpha x) = \alpha f(x)$.

Or $\forall x_1, x_2 \in V, \forall \alpha, \beta \in K : f(\alpha x_1 + \beta x_2) = \alpha f(x_1) + \beta f(x_2)$.

Example 1: The map $f : \mathbb{R}^2 \to \mathbb{R}^3$ defined as $f(x, y) = \left(x^2, x + y, 1\right)$ is not linear. We can easily find vectors for which the condition is false. For example:

$$f((1,0) + (0,0)) = f(1,0) = (1,1,1)$$

$$f(1,0) + f(0,0) = (1,1,1) + (0,0,1) = (1,1,2)$$

Hence, $f((1,0) + (0,0)) \neq f(1,0) + f(0,0)$

Example 2: $f : \mathbb{R}^2 \to \mathbb{R}^3$ defined as $f(x, y) = (3x - y, 0, 2y)$ is linear map:

(a) $f((x_1, y_1) + (x_2, y_2)) = f(x_1 + x_2, y_1 + y_1) = (3(x_1 + x_2), 0, 2(y_1 + y_2)) = (3x_1 - y_1, 0, 2y_1) + (3x_2 - y_2, 0, 2y_2) \Rightarrow f((x_1, y_1) + (x_2, y_2)) = f(x_1, y_1) + f(x_2, y_2)$.

(b) $f(\alpha.(x, y)) = f(\alpha.x, \alpha.y) = (3(\alpha x) - (\alpha y), 0, 2(\alpha.y)) = \alpha.(3x - y, 0, 2y) = \alpha f(x, y)$.

Properties. Here are some simple properties of linear maps $f : V \to W$.

(a) $f(0_V) = 0_W$.

(b) $f(-x) = -f(x)$.

(c) If V_1 is a subspace of V, then $f(V_1)$ is a subspace of W.

(d) If W_1 is a subspace of W, then $f^{-1}(W_1)$ is a subspace of V.

(e) The composite map of two linear maps is a linear map.

5.2.1 Linear maps and dimension

The kernel of a linear map

Definition. (**Kernel**)

The **kernel** (or **null space**) of such a f , denoted by $\ker f$, is the set of all x in V such that $f(x) = 0_W$ (the zero vector in W):

$$\ker f = \{x \in V, f(x) = 0_W\} = f^{-1}(\{0_W\})$$

The image of a linear map

Definition. (**Image**)

The **image** of f , denoted by Imf, is the set of all vectors in W of the form $f(x)$ for some x in V.

$$Imf = \{f(x), x \in V\} = f(V)$$

Properties. Let $f : V \to W$ be a linear map.

(a) The kernel of f is a subspace of V.

(b) The image of f is a subspace of W.

Proposition. Let $f : V \to W$ be a linear map.

(a) f is injective if and only if $\ker f = \{0_V\}$.

(b) f is surjective if and only if $Imf = W$.

Example: The map $f : \mathbb{R}^3 \to \mathbb{R}^2$ defined as $f(x, y, z) = (x + y, z)$ is not injective and surjective.

- f is injective $\Leftrightarrow \ker f = \{0_{\mathbb{R}^3}\}$.

$$\begin{aligned}
\ker f &= \left\{X \in \mathbb{R}^3, f(X) = (0,0)\right\} \\
&= \left\{(x,y,z) \in \mathbb{R}^3, f(x,y,z) = (0,0)\right\} \\
&= \left\{(x,y,z) \in \mathbb{R}^3, (x+y,z) = (0,0)\right\} \\
&= \left\{(x,y,z) \in \mathbb{R}^3, x+y=0 \text{ and } z=0\right\} \\
&= \left\{(x,y,z) \in \mathbb{R}^3, x=-y \text{ and } z=0\right\} \\
&= \{(-y,y,0), y \in \mathbb{R}\}
\end{aligned}$$

For example: $(-1,1,0) \in \ker f \Rightarrow \ker f \neq \{0_{\mathbb{R}^3}\}$. Hence f is not injective.

- f is surjective $\Leftrightarrow Imf = \left\{f(x), x \in \mathbb{R}^3\right\} = \mathbb{R}^2$.

$$\begin{aligned}
Imf &= \left\{f(X), X \in \mathbb{R}^3\right\} \\
&= \{(x+y,z) : x,y,z \in \mathbb{R}\} \\
&= \{x(1,0) + y(1,0) + z(0,1) : x,y,z \in \mathbb{R}\}
\end{aligned}$$

Hence, Imf is generated by two vectors $(1,0), (1,0)$ which are the *canonical basis* of $\mathbb{R}^2$. Then $Imf = \mathbb{R}^2$ and f is surjective.

Proposition. Let $f : V \to W$ be a linear map, with V finite-dimensional. Then:

$$f \text{ bijective} \Leftrightarrow f \text{ injective} \Leftrightarrow f \text{ surjective}$$

Proposition. Let $f : V \to W$ be a linear map, with V finite-dimensional. Then:

$$\dim V = \dim \ker f + \dim Imf$$

The rank of a linear map

Definition. (Rank)

The rank of a linear map f is the dimension of its image, written rank$f$:

$$\text{rank} f = \dim Imf$$

Example: Find $\ker f$, Imf and rank$f$ of the map $f : \mathbb{R}^4 \to \mathbb{R}^3$ defined as:

$$f(x, y, z, t) = (x - y, z + t, x - y + z)\,.$$

$$\begin{aligned}
\ker f &= \left\{X \in \mathbb{R}^4, f(X) = (0,0,0)\right\} \\
&= \left\{(x,y,z,t) \in \mathbb{R}^4, f(x,y,z,t) = (0,0,0)\right\} \\
&= \left\{(x,y,z,t) \in \mathbb{R}^4, (x-y, z+t, x-y+z) = (0,0)\right\} \\
&= \left\{(x,y,z,t) \in \mathbb{R}^4, x-y=0 \ \wedge \ z+t=0 \wedge x-y+z=0\right\} \\
&= \left\{(x,y,z) \in \mathbb{R}^3, \quad x=y \ \wedge \ z=t=0\right\} \\
&= \{(x,x,0,0)\,, \quad x \in \mathbb{R}\} = \{x.\,(1,1,0,0)\,, x \in \mathbb{R} \ \}
\end{aligned}$$

Hence , $\ker f = [\{(1,1,0,0)\}]$.

$$\begin{aligned}
Imf &= \{f(x,y,z,t)/ \ x,y,z,t \in \mathbb{R}\} \\
&= \{(x-y, z+t, x-y+z)\,/ \ x,y,z,t \in \mathbb{R}\} \\
&= \{(x-y)\,.\,(1,0,1) + t.\,(0,1,0) + z\,(0,1,1)\,/ \ x,y,z,t \in \mathbb{R}\}
\end{aligned}$$

Hence , $Imf = [\{(1,0,1)\,,(0,1,0)\,,(0,1,1)\}]$

Let $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$:

$$\alpha_1 (1,0,1) + \alpha_2 (0,1,0) + \alpha_3 (0,1,1) = (0,0,0) \quad \Rightarrow \quad (\alpha_1, \alpha_2 + \alpha_3, \alpha_1 + \alpha_3) = (0,0,0)\,.$$

$$\Rightarrow \quad \begin{cases} \alpha_1 = 0\,. \\ \alpha_2 + \alpha_3 = 0. \\ \alpha_1 + \alpha_3 = 0. \end{cases}$$

$$\Rightarrow \quad \alpha_1 = \alpha_2 = \alpha_3 = 0.$$

Therefore, the set $\{(1,0,1), (0,1,0), (0,1,1)\}$ is linearly independent and it is the basis of Imf.

$$\text{rank} f = \dim Imf = 3.$$

...

Appendix

Summary of the courses translated into arabic

Let’s REMEMBER translated

عمليات على النشر الحدود

إذا كانت f و g دالتان تقبلان نشرا محدودا جوار الصفر من الدرجة n فإن الدوال $f+g$, $(\alpha \in \mathbb{R}^*)$ $\alpha \cdot f$, $f \cdot g$, و f/g حيث $g(x) \neq 0$, هي دوال تقبل نشرا محدودا جوار الصفر من الدرجة n

بوضع:

$$f(x) = a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n + o(x^n)$$

$$g(x) = b_0 + b_1 x + b_2 x^2 + \cdots + b_n x^n + o(x^n)$$

ومنه:

- $(f+g)(x) = (a_0 + b_0) + (a_1 + b_1)x + (a_2 + b_2)x^2 + \cdots + (a_n + b_n)x^n + o(x^n)$
- $(\alpha \cdot f)(x) = (\alpha a_0) + (\alpha a_1)x + (\alpha a_2)x^2 + \cdots + (\alpha a_n)x^n + o(x^n)$
- $(f \cdot g)(x) = (a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n) \cdot (b_0 + b_1 x + b_2 x^2 + \cdots + b_n x^n) + o(x^n)$

  فيما يخص الجداء نحتفظ ماعدا بالحدود التي لها درجة أصغر من أو تساوي من الدرجة n .

- $\left(\frac{f}{g}\right)(x) = \frac{a_0 + a_1 x + a_2 x^2 + \cdots + a_n x^n}{b_0 + b_1 x + b_2 x^2 + \cdots + b_n x^n} + o(x^n)$

نقوم بالقسمة الاقليدية للبسط على المقام حسب القوى المتزايدة (الحد ذو أصغر درجة على الحد ذو أصغر درجة)

نشر مركب دالتين

إذا كانت g دالة تقبل نشرا محدودا جوار الصفر من الدرجة n و f دالة تقبل نشرا محدودا جوار $g(0)$ و $g(0) = 0$ فإن الدالة المركبة $f \circ g$ تقبل نشرا محدودا جوار الصفر من الدرجة n , بحيث يتم استبدال نشر g في النشر المحدود للدالة f و الاحتفاظ ماعدا بالحدود ذات الدرجة أقل أو تساوي من الدرجة n.

النشر المحدود جوار نقطة

نقول أن الدالة f تقبل نشرا محدودا **جوار النقطة** x_0 من الدرجة n إذا كانت F دالة تقبل نشرا محدودا **جوار الصفر** من الدرجة n حيث $F(X) = f(x) = f(X + x_0)$ حيث :

$$F(X) = a_0 + a_1 X + a_2 X^2 + \cdots + a_n X^n + o(X^n)$$

$$f(x) = a_0 + a_1(x - x_0) + a_2(x - x_0)^2 + \cdots + a_n(x - x_0)^n + o((x - x_0)^n)$$

النشر المحدود جوار المالانهاية ∞

نقول أن الدالة f تقبل نشرا محدودا **جوار** ∞ من الدرجة n إذا كانت F دالة تقبل نشرا محدودا **جوار** **الصفر** من الدرجة n حيث $F(X) = f(x) = f(\frac{1}{X})$ حيث :

$$F(X) = a_0 + a_1X + a_2X^2 + \cdots + a_nX^n + o(X^n)$$

$$f(x) = a_0 + \frac{a_1}{x} + \frac{a_2}{x^2} + \cdots + \frac{a_n}{x^n} + o\left(\frac{1}{x^n}\right)$$

تطبيق النشر المحدود في حساب النهايات

يستعمل النشر المحدود في إزالة حالات عدم التعيين بحيث يتم نشر التوابع جوار النقطة المطلوب حساب النهاية عندها وإلى أصغر درجة تمكننا من إزالة حالة عدم التعيين.

Appendix E

Vector Space and Linear Maps translated



الفضاء الشعاعي والتطبيقات الخطية

الفضاء الشعاعي

تحت كل فضاء شعاعي (سيتم تعريفه قريبًا) يوجد **حقل السلميات** K .

الحقل عبارة عن مجموعة من العناصر نعرف عليها العمليات الأساسية الأربع $(+,-,\times,\div)$ بخصائصها الاعتيادية (التبديل ، التجميع ، التوزيع).

أمثلة: $\mathbb{Q}, \mathbb{R}, \mathbb{C}$ لها بنية حقل ولكن $\mathbb{N}$ ليس لها بنية حقل (لا يمكن الطرح و القسمة دوما) $\mathbb{Z}$ ليس لها بنية حقل (لا يمكن القسمة دوما).

تعريف الفضاء الشعاعي

الفضاء الشعاعي على الحقل K هو مجموعة V غير خالية من العناصر , تسمى **أشعة** ، والتي يتم تعريف عمليتين عليها:

العملية الداخلية

$$+: \begin{array}{ccc} V \times V & \longrightarrow & V \\ (u,v) & \longmapsto & u+v \end{array}$$

العملية الخارجية

$$\cdot: \begin{array}{ccc} K \times V & \longrightarrow & V \\ (\alpha,u) & \longmapsto & \alpha u \end{array}$$

بحيث تتحقق الخواص التالية

1. $\forall u,v \in V:\ u+v=v+u$
2. $\forall u,v,w \in V:\ (u+v)+w=u+(v+w)$
3. $\forall u \in V, \exists 0_V \in V: u+0_V=u$ (0_V هو الشعاع المعدوم)
4. $\forall u \in V, \exists(-u) \in V:\ u+(-u)=0_V$
5. $\forall u,v \in V, \forall \alpha \in K: \alpha\,(u+v)=\alpha u+\alpha v$
6. $\forall u \in V, \forall \alpha,\ \beta \in K: (\alpha+\beta)u=\alpha u+\beta u$
7. $\forall u \in V, \forall \alpha,\ \beta \in K: (\alpha\,\beta)u=\alpha(\beta u)$
8. $\forall u \in V,\ 1\cdot u=u$

مثال: $\mathbb{Q}, \mathbb{C}$, $\mathbb{R}^3$, $\mathbb{R}^2$, $\mathbb{R}$ فضاءات شعاعية على $\mathbb{R}$

الفضاء الشعاعي الجزئي

تعريف1: **الفضاء الشعاعي الجزئي** من الفضاء V هو المجموعة الجزئية غير الخالية F من V تحقق الخاصيتين:

$$1.\ \forall u, v \in F:\ u + v \in F$$
$$2.\ \forall u \in F, \forall \alpha \in K: \alpha u \in F$$

تعريف2: **الفضاء الشعاعي الجزئي** من الفضاء V هو المجموعة الجزئية غير الخالية F من V تحقق الخاصية :

$$\forall u, v \in F, \forall \alpha, \beta \in K: \alpha u + \beta v \in F$$

نتيجة

- $\{0_V\}$ هو فضاء شعاعي جزئي من V .
- V هو فضاء شعاعي جزئي من نفسه.

مثال: $\mathbb{R}, \mathbb{R}^2, \mathbb{R}^3$ فضاءات شعاعية جزئية من $\mathbb{R}^3$

المزج الخطي

ليكن V **فضاء شعاعي** على الحقل K . نقول أن الشعاع u هو مزج خطي للأشعة $v_1, v_2, \dots, v_n$ من V إذا كان:

$$\exists \alpha_1, \alpha_2, \dots, \alpha_n \in K: u = \alpha_1 v_1 + \alpha_2 v_2 + \cdots + \alpha_n v_n$$

الاستقلال والارتباط الخطي

تعريف: لتكن جملة الأشعة $S = \{v_1, v_2, \dots, v_n\}$ من V .

نقول أن الجملة S **مستقلة خطيا** إذا كانت كل السلميات منعدمة من أجل كل مزج خطي منعدم لأشعة S أي:

$$\forall \alpha_1, \alpha_2, \dots, \alpha_n \in K: \alpha_1 v_1 + \alpha_2 v_2 + \cdots + \alpha_n v_n = 0_V \Rightarrow \alpha_1 = \alpha_2 = \cdots = \alpha_n = 0_K$$

واذا لم يتحقق هذا الشرط قلنا أن الجملة S **مرتبطة خطيا**

تعريف: لتكن جملة الأشعة $S = \{v_1, v_2, \dots, v_n\}$ من V

نقول أن الجملة S **مرتبطة خطيا** إذا وجدت سلميات غير منعدمة من أجل كل مزج خطي منعدم لأشعة S أي:

$$\exists \alpha_1, \alpha_2, \dots, \alpha_n \in K: \alpha_i \neq 0_K, i = 1, .., n \ \wedge \ \alpha_1 v_1 + \alpha_2 v_2 + \cdots + \alpha_n v_n = 0_V$$

نتيجة

- $\{u\}$ **مستقلة خطيا** $\Leftrightarrow$ $u \neq 0_V$
- S **مرتبطة خطيا** $\Rightarrow$ $0_V \in S = \{v_1, v_2, \dots, v_n\}$

الجملة المولدة

تعريف: ليكن V فضاء شعاعي و جملة منتهية من الأشعة $S = \{v_1, v_2, \dots, v_n\}$ من V, تسمى

جملة مولدة إذا كان كل شعاع $u \in V$ يكتب على شكل مزج خطي لأشعة S و نرمز بالرمز: $V = [S]$, معناه:

$$\forall u \in V, \exists \alpha_1, \alpha_2, \dots, \alpha_n \in K: u = \alpha_1 v_1 + \alpha_2 v_2 + \cdots + \alpha_n v_n$$

أساس الفضاء الشعاعي

تعريف: ليكن V فضاء شعاعي و جملة منتهية من الأشعة $S = \{v_1, v_2, \dots, v_n\}$ من V,

تسمى الجملة S **أساس** إذا وفقط إذا كان:

1. S **جملة مولدة** للفضاء V , أي $V = [S]$.
2. S **جملة مستقلة خطيا.**

مثال: $S = \{e_1 = (1,0), e_2 = (0,1)\}$ أساس للفضاء الشعاعي $\mathbb{R}^2$

خاصية: إذا كان S أساس للفضاء الشعاعي V فان **كل شعاع يكتب كمزج خطي وحيد** لأشعة S .

بعد الفضاء الشعاعي

تعريف: ليكن V فضاء شعاعي على الحقل K , إذا وجد أساس للفضاء V يحتوي على عدد منته n من الأشعة , يسمى العدد n **بعد منته للفضاء الشعاعي** V , ونكتب: $\dim V = n$

مثال1: $\{e_1 = (1,0,0), e_2 = (0,1,0), e_3 = (0,0,1)\}$ هو أساس للفضاء الشعاعي $\mathbb{R}^3$ على $\mathbb{R}$, ومنه: $\dim \mathbb{R}^3 = 3$.

مثال2: $\{1, i\}$ هو أساس للفضاء الشعاعي $\mathbb{C}$ على $\mathbb{R}$, $\dim \mathbb{C} = 2$.

خواص إذا كان V فضاء شعاعي ذو بعد منته n , إذن:

- كل جملة **مولدة** تحتوي على الأقل على n **شعاع** .
- جميع الأسس للفضاء الشعاعي V تحتوي على **نفس العدد من الأشعة**.
- فإذا كانت الجملة $\{v_1, v_2, \dots, v_n\}$ **مستقلة خطيا** فهي تشكل **أساسا** للفضاء V .
- فإذا كانت الجملة $\{v_1, v_2, \dots, v_n\}$ **مولدة** فهي تشكل **أساسا** للفضاء V .
- إذا كان F **فضاء شعاعي جزئي** من الفضاء V إذن: $\dim F \leq \dim V$ و إذا كان $\dim F = \dim V$ فان $F = V$.

التطبيق الخطي

تعريف: ليكن V, W فضائين شعاعين على نفس الحقل K , حيث:

$$f: \begin{array}{ccc} V & \longrightarrow & W \\ x & \longmapsto & f(x) \end{array}$$

نقول أن f هو **تطبيق خطي** من V نحو W إذا وفقط إذا كان:

$$\begin{cases} (1) & \forall x_1, x_2 \in V: \quad f(x_1 + x_2) = f(x_1) + f(x_2) \\ (2) & \forall \alpha \in K: \quad f(\alpha x) = \alpha f(x) \end{cases}$$

أو

$$\forall x_1, x_2 \in V, \forall \alpha, \beta \in K: f(\alpha x_1 + \beta x_2) = \alpha f(x_1) + \beta f(x_2)$$

خواص ليكن f **تطبيق خطي** من V نحو W

1. $f(0_V) = 0_W$
2. $f(-x) = -f(x)$
3. إذا كان V_1 **فضاء شعاعي جزئي** من الفضاء V فإن $f(V_1)$ **فضاء شعاعي جزئي** من الفضاء W
4. إذا كان W_1 **فضاء شعاعي جزئي** من الفضاء W فإن $f^{-1}(W_1)$ **فضاء شعاعي جزئي** من الفضاء V
5. مركب **تطبيقين خطيين** هو **تطبيق خطي**

التطبيقات الخطية والبعد

نواة تطبيق خطي

تعريف: نواة تطبيق خطي , ونرمز لها بالرمز $Ker\, f$, و هي مجموعة كل العناصر x من الفضاء V حيث $f(x) = 0_W$ ونكتب:

$$Ker\, f = \{x \in V, f(x) = 0_W\} = f^{-1}(\{0_W\})$$

صورة تطبيق خطي

تعريف: صورة تطبيق خطي , ونرمز لها بالرمز $Im\, f$, و هي مجموعة صور عناصر الفضاء V ونكتب:

$$Im\, f = \{f(x), x \in V\} = f(V)$$

خواص ليكن f تطبيق خطي من V نحو W

1. $Ker\, f$ هي فضاء شعاعي جزئي من الفضاء V.
2. $Im\, f$ هي فضاء شعاعي جزئي من الفضاء W.

التطبيق الخطي المتباين و الغامر

نظرية: ليكن f تطبيق خطي من V نحو W

1. f متباين إذا وفقط إذا كان $Ker\, f = \{0_V\}$
2. f غامر إذا وفقط إذا كان $Im\, f = W$

نظرية: ليكن f تطبيق خطي من V نحو W , مع العلم أن V ذو بعد منته. إذن:

$$\dim V = \dim Ker\, f + \dim Im\, f$$

التطبيق الخطي التقابلي

نظرية: ليكن f تطبيق خطي من V نحو W , مع العلم أن V ذو بعد منته. إذن:

- f تقابلي إذا وفقط إذا كان f متباين
- f تقابلي إذا وفقط إذا كان f غامر

رتبة التطبيق الخطي

تعريف: رتبة تطبيق خطي هي بعد صورته $Im\, f$, ونرمز لها بالرمز: $rank\, f$

ونكتب:

$$\boxed{rank\, f = \dim Im\, f}$$

Knowing is not enough,
We must APPLY
المعرفة لا تكفي ،
يجب أن نطبق
Willing is not enough,
We must DO
الرغبة لا تكفي ،
يجب أن نفعل
Bruce Lee

References

1. ركاب صورية، بوشقوف مسعودة وعرعار نورية . دروس الرياضيات 1-السنة الدراسية 2016-2018.

2. Guy Laffaille, Christian Pauly, Cours d'analyse 1, Licence 1er semestre, 2006.

3. Roberto's Notes on Differential Calculus, chapter 2: Resolving indeterminate forms, 2018.

4. Ion Goian, Raisa Grigor, Vasile Marin, Florentin Smarandache, ALGEBRAIC PROBLEMS AND EXERCISES FOR HIGH SCHOOL, The Educational Publisher, Colombus, 2015.

5. CHARLES C. PINTER , A Book of SET THEORY, Bucknell University, New York, 2014.

6. Abderezak Ould Houcine, Les Développements Limités, 2003-2004.

7. Calvin Jongsma, Discrete Mathematics: Chapter 6, Functions & Equivalence Relations, 2016.

8. James Tanton, Ph.D, ENCYCLOPEDIA OF Mathematics, New York, 2005.

9. Sigurd Angenent, MATH 221 , FIRST SEMESTER, CALCULUS, 2010 ,

10. Ronald van Luijk, Linear Algebra I, 2012.

11. AMIN WITNO, LOGIC& SET THEORY, 2014.

12. Johannes Kepler, Mathematics for Chemists 2, Lecture 6: Linear algebra I, 2012.

13. Sean Mauch, Introduction to Methods of Applied Mathematics or Advanced Mathematical Methods for Scientists and Engineers, 2003.

14. Milos Hauskrecht , CS 441 Discrete Mathematics for CS Lecture 7. Spring 2018.

15. Robert Howlett, Vector Space Theory, A course for second year students, The university of Sydney, Math 2902, 2001.

16. Jérom Dubois, "Algébre et Analyse élèmentaires", 2013-2014.

17. Chapter Ten APPROXIMATING FUNCTIONS, 446-476, math.arizona.

18. Mathematics and Statistics – I, Mathematical Logic, Target Publications- Transforming Lives through Learning. published by the Maharashtra State Board of Secondary and Higher Secondary Education, Pune, 2013.

19. Gloria Faccanoni , M11 Mathématiques, Recueil d'exercices corrigés et aide-mémoire, Année 2014 – 2015.

20. Mathematics in the Islamic World (8th – 15th Century), Encyclopædia Britannica.

MATHEMATICS I

1. Logic and mathematical reasoning.
2. Sets, Relations and Applications.
3. Real-valued functions of a real variable.
4. Finite Expansions.
5. Vector Space and Linear Maps.

KESMIA Mounira